

# Fetal Alcohol Syndrome and Fetal Alcohol Spectrum Disorders

LEEANNE DENNY, MD; SARAH COLES, MD; and ROBIN BLITZ, MD  
*University of Arizona College of Medicine, Phoenix, Arizona*

Fetal alcohol syndrome (FAS) and fetal alcohol spectrum disorders (FASD) result from intrauterine exposure to alcohol and are the most common nonheritable causes of intellectual disability. The percentage of women who drink or binge drink during pregnancy has increased since 2012. FAS is commonly missed or misdiagnosed, preventing affected children from receiving needed services in a timely fashion. Diagnosis is based on the presence of the following clinical features, all of which must be present: prenatal and/or postnatal growth retardation, facial dysmorphism, central nervous system dysfunction, and neurobehavioral disabilities. FASD is a broader diagnosis that encompasses patients with FAS and others who are affected by prenatal alcohol exposure but do not meet the full criteria for FAS. Management is multidisciplinary and includes managing comorbid conditions, providing nutritional support, managing behavioral problems and educational difficulties, referring patients for rehabilitative therapies, and educating parents. The Centers for Disease Control and Prevention and other organizations recognize no safe amount of alcohol consumption during pregnancy and recommend complete abstinence from alcohol. All women should be screened for alcohol use during preconception counseling and prenatal care, and alcohol use should be addressed with brief interventions. (*Am Fam Physician*. 2017;96(8):515-522. Copyright © 2017 American Academy of Family Physicians.)



More online  
at <http://www.aafp.org/afp>.

**CME** This clinical content conforms to AAFP criteria for continuing medical education (CME). See CME Quiz on page 498. Author disclosure: No relevant financial affiliations.

► **Patient information:** A handout on this topic is available at <https://familydoctor.org/condition/fetal-alcohol-syndrome>.

**F**etal alcohol spectrum disorders (FASD) result from prenatal exposure to alcohol and include fetal alcohol syndrome (FAS), partial fetal alcohol syndrome (PFAS), alcohol-related neurodevelopmental disorder, and alcohol-related birth defects.<sup>1</sup> FAS is the most severe form of FASD.<sup>2</sup>

According to the Centers for Disease Control and Prevention, the percentage of pregnant women who consume alcohol increased from 7.6% in 2012 to 10.2% in 2015, and the number of pregnant women reporting binge drinking (four or more alcoholic beverages at once) increased from 1.4% to 3.1%.<sup>3,4</sup> These trends are concerning because alcohol is the most common teratogen, and FASD is the most common cause of nonheritable

intellectual disability.<sup>5</sup> Binge drinking is associated with the development of behavioral problems and physical deformities.<sup>6</sup>

Although there is wide variation in the estimated prevalence of FAS/FASD, FAS is thought to occur in 0.3 to 0.8 per 1,000 children in the United States and in 2.9 per 1,000 globally.<sup>7,8</sup> The prevalence of FASD is estimated at 33.5 per 1,000 children in the United States and 22.8 per 1,000 globally.<sup>8</sup> In the United States, FASD is least prevalent in Hispanic children and most prevalent in Native Americans and Alaska Natives.<sup>4</sup> FAS is diagnosed at an average age of 48.3 months<sup>9</sup>; however, it is commonly missed or misdiagnosed, preventing affected children from receiving needed services in a timely fashion.

FASD carries a significant economic burden. Children with FAS who are enrolled in Medicaid have annual mean medical expenses nine times higher than those for children without FAS, equating to a median annual expenditure of \$6,670 per child (vs. \$518 for those without FAS).<sup>10</sup>

## Diagnosis

Any child who was exposed to alcohol prenatally or presents with growth retardation, facial dysmorphism, central nervous system

### WHAT IS NEW ON THIS TOPIC: FETAL ALCOHOL SPECTRUM DISORDERS

According to the Centers for Disease Control and Prevention, the percentage of pregnant women who consume alcohol increased from 7.6% in 2012 to 10.2% in 2015, and the number of pregnant women reporting binge drinking (at least four alcoholic beverages at once) increased from 1.4% to 3.1%.

A study demonstrated that more than one-half of children with fetal alcohol spectrum disorders do not consume the recommended dietary allowance of fiber, calcium, or vitamins D, E, and K.

## Fetal Alcohol Syndrome

dysfunction, or neurobehavioral disabilities—the key manifestations of FASD—should prompt consideration of FASD.<sup>11</sup> The assessment and diagnosis require a multidisciplinary team (Table 1<sup>1,12</sup>) and should include neuropsychological assessment.<sup>1</sup>

Diagnosis begins with assessment of prenatal alcohol exposure, including quantity of alcohol consumed per occasion, frequency of use, and timing of consumption during pregnancy. Prenatal alcohol exposure is defined as at least one of the following documented findings: (1) six or more drinks per week for two or more weeks during pregnancy; (2) three or more drinks per occasion on two or more occasions during pregnancy; (3) alcohol-related social or legal problems around the time of pregnancy; (4) intoxication during pregnancy documented by blood, breath, or urinary alcohol testing; (5) positive test for alcohol exposure biomarkers during pregnancy (fatty acid ethyl esters, phosphatidylethanol, and ethyl glucuronide in maternal hair, fingernails, urine, or blood, or in placenta or meconium); (6) increased prenatal risk associated with alcohol use during pregnancy as assessed by a validated screening tool. Documentation includes drinking levels reported by the mother three months before pregnancy recognition or at the time of

**Table 1. Multidisciplinary Team for the Care of Patients with Fetal Alcohol Spectrum Disorders**

Audiologist	Physical therapist
Cardiologist	Play therapist
Developmental pediatrician	Primary care physician
Developmental therapist	Psychiatrist
Family therapist	Psychotherapist
Nephrologist	Sensory integration therapist
Neurologist	Social worker
Occupational therapist	Special education teachers
Ophthalmologist	Speech-language pathologist

NOTE: Although not all children with fetal alcohol spectrum disorders will require care from all disciplines listed, referrals should be initiated based on co-occurring conditions and needs.

Information from references 1 and 12.

a positive pregnancy test. Information must be obtained by the mother or a reliable source, such as family member, social service agency, or medical record.<sup>1</sup>

Exposure to other teratogens should also be assessed, because women who consume alcohol during pregnancy are more likely to use other drugs.<sup>1</sup> The diagnostic criteria for FAS or PFAS do not require confirmed alcohol use if characteristic findings are present.<sup>1,11</sup> However, a confirmed absence of alcohol exposure rules out the diagnoses. Confirmation of alcohol exposure is required

**Table 2. Diagnosis of Fetal Alcohol Spectrum Disorders**

Documented prenatal alcohol exposure	Facial dysmorphism <sup>*</sup>	Growth deficiency <sup>†</sup>	Central nervous system dysfunction <sup>‡</sup>	Neurobehavioral impairment <sup>§</sup>	Diagnosis
+	+	+	+	+	Fetal alcohol syndrome
–	+	+	+	+	Fetal alcohol syndrome
+	+	+	–	+	Partial fetal alcohol syndrome
+	+	–	+	+	Partial fetal alcohol syndrome
+	+	–	–	+	Partial fetal alcohol syndrome
–	+	+	–	+	Partial fetal alcohol syndrome
–	+	–	+	+	Partial fetal alcohol syndrome
+	–	–	–	+	Alcohol-related neurodevelopmental disorder¶

\*—Requires two or more of the following: short palpebral fissure, thin vermilion border of the upper lip, and smooth philtrum.

†—May be prenatal or postnatal and includes height and/or weight  $\leq$  10th percentile on appropriate growth curve.


‡—Must include one of the following: head circumference  $\leq$  10th percentile on appropriate growth curve, structural brain abnormalities, or recurrent nonfebrile seizures with no other identifiable cause.

§—Requires evidence of global impairment or deficit in at least one neurobehavioral domain  $\geq$  1.5 standard deviations below mean.

||—Requires evidence of global impairment or deficit in at least two neurobehavioral domains.

¶—Cannot be definitively diagnosed in children younger than three years.

Information from reference 1.



**Figure 1.** Facial features associated with fetal alcohol spectrum disorders.

Reprinted with permission from Wattendorf DJ, Muenke M. *Fetal alcohol spectrum disorders*. *Am Fam Physician*. 2005;72(2):279.


for diagnosis of alcohol-related neurodevelopmental disorder and alcohol-related birth defects.<sup>1</sup>

#### KEY DIAGNOSTIC CRITERIA

As previously noted, FASD comprises four distinct categories: FAS, PFAS, alcohol-related neurodevelopmental disorder, and alcohol-related birth defects. Each category is distinguished by the presence or absence of characteristic facial dysmorphology, growth retardation, central nervous system dysfunction, and neurobehavioral disabilities (Table 2).<sup>1</sup>

Characteristic facial dysmorphology associated with FASD includes short palpebral fissures (10th percentile or less for age and racial norms), a thin vermilion border of the upper lip, and a smooth philtrum<sup>1</sup> (Figure 1<sup>13</sup>). Two of the three characteristic features are required for the diagnosis of FAS or PFAS. Palpebral fissures can be measured using a small plastic ruler, noting the distance between the endocanthion (where the eyelids meet medially) and exocanthion (where they meet laterally). The ruler should be angled to follow the curve of the zygoma.<sup>1</sup> The presence of a thin vermilion border and smooth philtrum is scored objectively using the lip-philtrum scoring guide (Figure 2).<sup>14</sup> Scores of 4 or 5 are consistent with FAS or PFAS.

Growth retardation is defined as the 10th percentile or less using height and weight measurements on standard growth curves.<sup>1</sup> For central nervous system dysfunction to qualify as consistent with FASD, it must include deficient brain growth, abnormal structure, or abnormal neurophysiology. This can be documented as a head circumference in the 10th percentile or less on appropriate growth curves, structural brain abnormalities, or recurrent nonfebrile seizures with no other



**Figure 2.** Lip-Philtrum Guide I is used to rank upper lip thinness and philtrum smoothness. Ranks 4 and 5 reflect the thin lip and smooth philtrum that characterize the FAS facial phenotype. Rank 3 represents the general population mean.

Reprinted with permission from FAS Diagnostic & Prevention Network. Copyright 2017, Susan Ashley PhD, University of Washington. <http://depts.washington.edu/fasdpn/htmls/lip-philtrum-guides.htm>.

identifiable cause.<sup>1</sup> Magnetic resonance imaging has identified structural brain abnormalities in children with FASD (e.g., temporal lobe asymmetry, change in size or shape of corpus callosum, cerebellum, or basal ganglia), and it may be used in the evaluation of sus-

**Table 3. Conditions Commonly Occurring with Fetal Alcohol Spectrum Disorders**

System	Condition
Auditory	Chronic serous otitis media, conductive and/or neurosensory hearing loss
Cardiac	Aberrant great vessels, atrial septal defects, ventricular septal defects
Gastrointestinal	Enteric neuropathy
Musculoskeletal	Camptodactyly, clinodactyly (Figure 3), flexion contractures, hypoplastic nails, radioulnar synostosis, scoliosis, spinal malformations
Neurologic	Microcephaly, seizure disorder, spinal cord abnormalities, structural brain abnormalities (including corpus callosum, cerebellum, caudate, and hippocampus)
Ophthalmologic	Ptosis, retinal malformation, strabismus, visual impairment
Orofacial	Cleft lip, cleft palate
Psychiatric/ neuro- behavioral	Attention-deficit/hyperactivity disorder, conduct disorder, intellectual disability, language disorders, learning disabilities, mood disorders, oppositional defiant disorder, substance use disorders
Renal	Aplastic/dysplastic/hypoplastic kidneys, horseshoe kidney, hydronephrosis, ureteral duplications

Information from references 15, and 18 through 21.

involve a chromosomal microarray, cranial neuroimaging, and cardiac/abdominal ultrasonography.<sup>2</sup>

### Management

There is no cure for FASD.<sup>5</sup> There is a lack of evidence on which to base recommendations for optimal management; therefore, much of the management is based on expert opinion. Treatment consists of providing a medical home for the patient and family, managing comorbid conditions, providing nutritional support, addressing behavioral and emotional problems, arranging referrals for rehabilitative therapies (therapeutic intervention for those who have never developed a specific skill), coordinating care with a multidisciplinary team, and educating parents (Table 5). Early intervention is necessary to optimize health outcomes.<sup>11,29</sup>

pected FASD; it can also be helpful if there is a question about the differential diagnosis.<sup>1,15-17</sup>

Neurobehavioral disabilities in FASD include deficient global intellectual ability and cognition, and poor behavior, self-regulation, and adaptive skills. These domains should be measured using standardized testing, which often cannot be administered until after three years of age. A deficiency on these tests is characterized by scores of at least 1.5 standard deviations below the mean.<sup>1</sup> Alcohol-related neurodevelopmental disorder is diagnosed with documented prenatal alcohol exposure and neurobehavioral impairment in at least two domains in the absence of other defining characteristics for FAS.

Although they are not included in the diagnostic criteria for FAS or PFAS, multiple congenital abnormalities associated with prenatal alcohol exposure have been described for nearly every organ system (Table 3).<sup>15,18-21</sup> In the absence of defining criteria for FAS or PFAS, documented prenatal alcohol exposure and the presence of one or more major malformations known to result from prenatal alcohol exposure are diagnostic for alcohol-related birth defects<sup>1</sup> (eTable A, Figure 3<sup>13</sup>).

### Differential Diagnosis

The differential diagnosis for FASD includes a variety of chromosomal abnormalities, exposure to other teratogens, and behavioral and psychiatric diagnoses (Table 4).<sup>2,22-28</sup> If the diagnosis is uncertain, the workup should include referral to a developmental pediatrician or geneticist for further evaluation, which may



**Figure 3.** Hand features associated with fetal alcohol spectrum disorders include clinodactyly (curved fifth digit) and “hockey stick” crease (distal palmar crease widens between the second and third digits).

Reprinted with permission from Wattendorf DJ, Muenke M. Fetal alcohol spectrum disorders. *Am Fam Physician*. 2005;72(2):281.

**Table 4. Differential Diagnosis of Fetal Alcohol Spectrum Disorders**

<i>Condition</i>	<i>Cause</i>	<i>Features similar to fetal alcohol syndrome</i>	<i>Distinguishing features from fetal alcohol syndrome</i>
Aarskog syndrome	X-linked recessive, often mutations in <i>FGD1</i> , although others unknown	Broad philtrum, intellectual and neurobehavioral disabilities, small nose with anteverted nares, wide-spaced eyes	Brachydactyly, crease below lower lip, dental eruption problems, downward-slanting palpebral fissures, shawl scrotum (scrotum folds around penis), short stature that resolves with puberty, widow's peak
Bloom syndrome	Autosomal recessive chromosomal instability caused by mutation in <i>BLM</i>	Short stature with mild microcephaly, variably impaired intellectual ability	Café au lait spots; facial telangiectasia erythema; keel-shaped face; predisposition to early cancer, infertility, and immunodeficiency; sparse subcutaneous adipose tissue
Cornelia de Lange (Brachmann-de Lange) syndrome	Autosomal dominant from spontaneous mutations in <i>NIPBL</i> , <i>RAD21</i> , and <i>SMC3</i> , or X-linked dominant with mutations in <i>HDAC8</i> or <i>SMC1A</i>	Anteverted nares, depressed nasal bridge, growth impairment, hearing loss, intellectual disability, microcephaly, short stature, smooth philtrum, thin vermilion border	Arched eyebrows that meet in the middle (synophrys), downturned mouth, high arched palate, hypertrichosis, long eyelashes, short limbs
Dubowitz syndrome	Unknown; suspected autosomal recessive	Neurobehavioral disabilities (hyperactivity, impulsivity, and inattentiveness), epicanthal folds, intellectual disability, microcephaly, short palpebral fissures, short stature, wide-spaced eyes	Broad nasal tip, cryptorchidism, eczema-like skin disorder, high-pitched voice, shallow supraorbital ridge with nasal bridge near level of forehead
Fetal hydantoin syndrome	Prenatal exposure to phenytoin (Dilantin)	Depressed nasal bridge, growth deficits, occasional intellectual disability, wide-spaced eyes	Genitourinary defects, hirsutism, hypoplastic fingertips, low hairline, orofacial clefts, short neck, short nose with bowed upper lip
Fetal valproate syndrome	Prenatal exposure to valproate (Depacon)	Anteverted nares, epicanthal folds, long philtrum, thin vermilion border, wide-spaced eyes	Cardiac malformations, high forehead, infraorbital crease, neural tube defects, small mouth
Noonan syndrome	Autosomal dominant, often mutation in <i>PTPN11</i>	Epicanthal folds, intellectual disability, low nasal bridge, short stature, wide-spaced eyes	Bleeding diathesis, cryptorchidism, downward-slanting palpebral fissures, hypertrophic cardiomyopathy, keratoconus, low posterior hairline, pectus excavatum, protruding upper lip, pulmonary stenosis, webbed neck, wide mouth
Phenylalanine embryopathy	Maternal phenylketonuria	Epicanthal folds, growth impairment, intellectual disability, long philtrum, microcephaly, short palpebral fissures, small nose with anteverted nares, thin vermilion border	Cardiac malformations, hypertonia, prominent glabella, round facies
Toluene embryopathy	Prenatal exposure to toluene	Growth deficits, midface hypoplasia, short palpebral fissures, smooth philtrum, thin vermilion border	Bifrontal narrowing of the skull, downturned mouth, ear abnormalities, hair pattern abnormalities, large anterior fontanelle, micrognathia
Velocardiofacial syndrome	Autosomal dominant with microdeletion in chromosome 22q11	Intellectual disabilities, psychiatric disorders, small palpebral fissures	Cardiac malformations, cleft palate, long face with prominent nose, transient neonatal hypocalcemia, weak pharyngeal muscles resulting in hypernasal speech
Williams syndrome	Heterozygous 7q11.23 deletion, including elastin gene	Anteverted nares, depressed nasal bridge, epicanthal folds, growth impairment, intellectual disability, long philtrum, short nose, short palpebral fissures	Aortic and pulmonary stenosis, connective tissue disorders, endocrine abnormalities, full lips, hoarse voice, hypertension, periorbital fullness, poor to near-normal language skills, renal abnormalities, stellate pattern of iris, systemic arterial stenosis, wide mouth

Information from references 2, and 22 through 28.

## SORT: KEY RECOMMENDATIONS FOR PRACTICE

<i>Clinical recommendation</i>	<i>Evidence rating</i>	<i>References</i>
The diagnosis of fetal alcohol syndrome and partial fetal alcohol syndrome is based on defined clinical characteristics and does not require confirmed alcohol use during pregnancy.	C	1
Neurobehavioral testing should be conducted in all children with suspected fetal alcohol spectrum disorders when feasible. Comprehensive evaluation may not be possible using conventional assessment tools until after three years of age.	C	1
Contraception should be offered to women of childbearing age who drink. If they desire pregnancy, abstinence from alcohol should be recommended.	C	44
Pregnant women should be screened for alcohol use. This can be done using the TACER-3 tool.	C	42, 45, 46

*A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <http://www.aafp.org/afpsort>.*

### MANAGING COMORBID CONDITIONS

Children with FASD can have a range of comorbid conditions (*Table 3*)<sup>15,18-21</sup>; referrals to members of the multidisciplinary team are based on the specific needs identified. Because hearing and vision impairments are correlated with prenatal alcohol exposure, all children with suspected FASD should have hearing and vision screening.<sup>30,31</sup>

### NUTRITIONAL SUPPORT

Children with FASD are nutritionally and socially vulnerable and may benefit from nutritional education and support. By midchildhood, most of these children have spent, on average, one-fourth of their life with unmet basic needs and one-third of their life with someone who abuses alcohol or drugs.<sup>29</sup> One study showed that more than 50% of children with FASD do not consume the recommended dietary allowance of fiber, calcium, or vitamins D, E, and K.<sup>32</sup> It is important to regularly assess the child's height, weight, and body mass index and refer for support (e.g., nutritionist, social worker) when nutritional problems are identified.<sup>33</sup> Some children will require high-calorie foods and supplements.

### MANAGING BEHAVIORAL PROBLEMS

Children with FASD should be monitored and screened for behavioral problems. They have an increased risk

of attention-deficit/hyperactivity disorder (40% to 95%),<sup>34,35</sup> mood disorders (50%),<sup>36</sup> and oppositional defiant disorder (38%).<sup>35</sup> Medications can improve hyperactivity and impulsivity, but not symptoms of inattention.<sup>37,38</sup> Children with FASD and attention-deficit/hyperactivity disorder or other disruptive behaviors should be referred to a developmental pediatrician, psychologist, and/or psychiatrist. Behavioral interventions such as play therapy, children's friendship training, and specially trained case managers have reasonable evidence of effectiveness, but these resources have variable availability.<sup>37</sup>

### FAMILY SUPPORT

Children with FASD are at increased risk of physical and sexual violence, with 61% experiencing physical or sexual abuse or witnessing domestic violence by 12 years of age.<sup>29,39</sup> Sexual abuse should be considered in children who present with inappropriate sexual behaviors. Children with FASD who remain in the care of their biologic mother are more likely to experience family dysfunction and instability (e.g., divorce, unemployment, drug and alcohol use).<sup>25,29</sup> Those who are raised in stable homes have improved outcomes and are less likely to be expelled from or drop out of school, be arrested, or develop substance use problems.<sup>29</sup> Interventions should be aimed at stabilizing the home environment and improving parent-child interactions.<sup>11</sup> Such interventions include parental substance abuse referrals, child discipline courses, parent support groups, and child protective services.

### Prognosis

Prognosis varies with the degree of impairment. Persons with FASD are more likely to require special education, receive disability pensions, and be unemployed.<sup>40</sup> Those who receive early diagnosis and intervention (before 12 years of age) have significantly better outcomes, including a two- to fourfold reduction in rates of imprisonment and substance abuse.<sup>29</sup>

**Table 5. Patient Resources for Fetal Alcohol Spectrum Disorders**

**American Academy of Pediatrics Fetal Alcohol Spectrum Disorders Program**

<http://www.aap.org/en-us/advocacy-and-policy/aap-health-initiatives/fetal-alcohol-spectrum-disorders-toolkit/>

**Centers for Disease Control and Prevention**

<https://www.cdc.gov/ncbddd/fasd/> and <https://www.cdc.gov/ncbddd/fasd/alcohol-use.html>

**National Organization on Fetal Alcohol Syndrome (also contains resources for teachers)**

<http://www.nofas.org/parents/>

**Table 6. TACER-3 Screening Tool for Alcohol Misuse**

Component	Positive reply	Score	Question
Tolerance	≥ 2 drinks*	2	How many drinks does it take to make you feel high?
Annoyance	Yes	1	Has anybody ever annoyed you by complaining about your drinking?
Cut down	Yes	1	Have you ever felt you ought to cut down on your drinking?
Eye-opener	Yes	1	Have you ever needed a drink first thing in the morning to get going?

NOTE: A positive screening result is a score of 3 or more.

\*—One drink is the equivalent of 0.5 oz of absolute alcohol (approximately 12 oz of regular beer, 1.5 oz of liquor, or 4 oz of wine).

Adapted with permission from Chiodo LM, Delaney-Black V, Sokol RJ, Janisse J, Pardo Y, Hannigan JH. Increased cut-point of the TACER-3 screen reduces false positives without losing sensitivity in predicting risk alcohol drinking in pregnancy. *Alcohol Clin Exp Res.* 2014;38(5):1403.

## Prevention

The Centers for Disease Control and Prevention, the American Academy of Family Physicians, the American Academy of Pediatrics, and the American Congress of Obstetricians and Gynecologists recognize no safe amount of alcohol consumption during pregnancy and recommend complete abstinence.<sup>26,41-43</sup> Although many women abstain from alcohol when they learn they are pregnant, some consume alcohol before they find out. Contraception should be offered to women of childbearing age who drink; if they desire pregnancy, abstinence from alcohol should be recommended.<sup>44</sup> The American Congress of Obstetricians and Gynecologists recommends screening women in the first trimester for alcohol use, and Canadian guidelines recommend screening all pregnant women for alcohol use.<sup>42,45</sup> A useful screening tool is the TACER-3, which identifies women whose drinking may put their fetus at risk of FASD (Table 6).<sup>46</sup>

If alcohol use in pregnancy is identified, physicians should recommend cessation and offer group-based interventions such as Alcoholics Anonymous and alcohol rehabilitation centers.<sup>47</sup> Brief interventions that include the patient's partner improve FASD-related birth outcomes and should include assessing maternal understanding of healthy pregnancy behaviors, assisting the mother in setting the goal of abstinence from alcohol, planning alternative behaviors for when the temptation to drink arises, and inviting the partner to find methods to support the mother's abstinence from alcohol.<sup>48,49</sup>

This article updates a previous article on this topic by Wattendorf and Muenke.<sup>13</sup>

**Data Sources:** Sources searched include PubMed (OVID), Evidence Summary from the *AFP's* editors, Essential Evidence Plus, Cochrane database, and the Agency for Healthcare Research and Quality. Search terms included: fetal alcohol syndrome, fetal alcohol spectrum disorder, alco-

hol-related birth defects, maternal alcohol consumption, prenatal alcohol exposure. Search dates: February 2016, April 2016, May 2016, June 2016, July 2016, November 2016, and December 2016.

Figures 1 and 3 courtesy of Darryl Leja, National Human Genome Research Institute, National Institutes of Health, Bethesda, Md.

## The Authors

LEEANNE DENNY, MD, is residency faculty at the University of Arizona College of Medicine Family Medicine Residency, Phoenix.

SARAH COLES, MD, is residency faculty at the University of Arizona College of Medicine Family Medicine Residency.

ROBIN BLITZ, MD, is a clinical associate professor of child health at the University of Arizona College of Medicine, Phoenix.

Address correspondence to LeeAnne Denny, MD, Banner University Medical Center—Phoenix, 1300 N. 12th St., Phoenix, AZ 85006 (e-mail: LeeAnne.Denny@bannerhealth.com). Reprints are not available from the authors.

## REFERENCES

- Hoyme HE, Kalberg WO, Elliott AJ, et al. Updated clinical guidelines for diagnosing fetal alcohol spectrum disorders. *Pediatrics.* 2016;138(2):e20154256.
- de Sanctis L, Memo L, Pichini S, Tarani L, Vagnarelli F. Fetal alcohol syndrome: new perspectives for an ancient and underestimated problem. *J Matern Fetal Neonatal Med.* 2011;24(suppl 1):34-37.
- Tan CH, Denny CH, Cheal NE, Sniezek JE, Kanny D. Alcohol use and binge drinking among women of childbearing age—United States, 2011-2013. *MMWR Morb Mortal Wkly Rep.* 2015;64(37):1042-1046.
- Centers for Disease Control and Prevention (CDC). Alcohol use and binge drinking among women of childbearing age—United States, 2006-2010. *MMWR Morb Mortal Wkly Rep.* 2012;61(28):534-538.
- Joya X, Garcia-Algar O, Salat-Batlle J, Pujades C, Vall O. Advances in the development of novel antioxidant therapies as an approach for fetal alcohol syndrome prevention. *Birth Defects Res A Clin Mol Teratol.* 2015;103(3):163-177.
- Alvik A, Aalen OO, Lindemann R. Early fetal binge alcohol exposure predicts high behavioral symptom scores in 5.5-year-old children. *Alcohol Clin Exp Res.* 2013;37(11):1954-1962.
- Fox DJ, Pettygrove S, Cunniff C, et al.; Centers for Disease Control and Prevention (CDC). Fetal alcohol syndrome among children aged 7-9 years—Arizona, Colorado, and New York, 2010. *MMWR Morb Mortal Wkly Rep.* 2015;64(3):54-57.
- Roozen S, Peters GJ, Kok G, Townend D, Nijhuis J, Curfs L. Worldwide prevalence of fetal alcohol spectrum disorders: a systematic literature review including meta-analysis [published correction appears in *Alcohol Clin Exp Res.* 2016;40(7):1587]. *Alcohol Clin Exp Res.* 2016;40(1):18-32.
- Moberg DP, Bowser J, Burd L, Elliott AJ, Punyko J, Wilton G; Fetal Alcohol Syndrome Surveillance Program-FASSLink Team. Fetal alcohol syndrome surveillance: age of syndrome manifestation in case ascertainment. *Birth Defects Res.* 2014;100(9):663-669.
- Amendah DD, Grosse SD, Bertrand J. Medical expenditures of children in the United States with fetal alcohol syndrome. *Neurotoxicol Teratol.* 2011;33(2):322-324.
- Bertrand J, Floyd LL, Weber MK; Fetal Alcohol Syndrome Prevention Team, Division of Birth Defects and Developmental Disabilities, National

## Fetal Alcohol Syndrome

- Center on Birth Defects and Developmental Disabilities, Centers for Disease Control and Prevention (CDC). Guidelines for identifying and referring persons with fetal alcohol syndrome [published correction appears in *MMWR Morb Mortal Wkly Rep*. 2006;55(20):568]. *MMWR Recomm Rep*. 2005;54(RR-11):1-14.
12. Chasnoff IJ, Wells AM, King L. Misdiagnosis and missed diagnoses in foster and adopted children with prenatal alcohol exposure. *Pediatrics*. 2015;135(2):264-270.
  13. Wattendorf DJ, Muenke M. Fetal alcohol spectrum disorders. *Am Fam Physician*. 2005;72(2):279-282, 285.
  14. FAS Diagnostic & Prevention Network. Lip-philtrum guides. <http://depts.washington.edu/fasdpn/htmls/lip-philtrum-guides.htm>. Accessed January 30, 2017.
  15. Norman AL, Crocker N, Mattson SN, Riley EP. Neuroimaging and fetal alcohol spectrum disorders. *Dev Disabil Res Rev*. 2009;15(3):209-217.
  16. Stoos C, Nelsen L, Schissler KA, Elliot AJ, Kinney HC. Fetal alcohol syndrome and secondary schizophrenia: a unique neuropathological study. *J Child Neurol*. 2015;30(5):601-605.
  17. Cook JL, Green CR, Lilley CM, et al.; Canada Fetal Alcohol Spectrum Disorder Research Network. Fetal alcohol spectrum disorder: a guideline for diagnosis across the lifespan. *CMAJ*. 2016;188(3):191-197.
  18. Popova S, Lange S, Shield K, et al. Comorbidity of fetal alcohol spectrum disorder: a systematic review and meta-analysis. *Lancet*. 2016;387(10022):978-987.
  19. American Academy of Pediatrics Committee on Substance Abuse and Committee on Children with Disabilities. Fetal alcohol syndrome and alcohol-related neurodevelopmental disorders. *Pediatrics*. 2000;106(2 pt 1):358-361.
  20. Caputo C, Wood E, Jabbour L. Impact of fetal alcohol exposure on body systems: a systematic review. *Birth Defects Res C Embryo Today*. 2016;108(2):174-180.
  21. Gummel K, Ygge J. Ophthalmologic findings in Russian children with fetal alcohol syndrome. *Eur J Ophthalmol*. 2013;23(6):823-830.
  22. Douzgon S, Breen C, Crow YJ, et al. Diagnosing fetal alcohol syndrome: new insights from newer genetic technologies. *Arch Dis Child*. 2012;97(9):812-817.
  23. Burd L, Cotsonas-Hassler TM, Martsolf JT, Kerbeshian J. Recognition and management of fetal alcohol syndrome. *Neurotoxicol Teratol*. 2003;25(6):681-688.
  24. Jones KL. Fetal alcohol syndrome. In: *Smith's Recognizable Patterns of Human Malformation*. 6th ed. Philadelphia, Pa.: Elsevier Saunders; 2006:646.
  25. Toutain S, Lejeune C. Family management of infants with fetal alcohol syndrome or fetal alcohol spectrum disorders. *J Dev Phys Disabil*. 2008;20(5):425-436.
  26. National Center on Birth Defects and Developmental Disabilities; Centers for Disease Control and Prevention; U.S. Department of Health and Human Services; National Task Force on Fetal Alcohol Syndrome and Fetal Alcohol Effect. Fetal alcohol syndrome: guidelines for referral and diagnosis. July 2004. [http://www.cdc.gov/ncbddd/fasd/documents/FAS\\_guidelines\\_accessible.pdf](http://www.cdc.gov/ncbddd/fasd/documents/FAS_guidelines_accessible.pdf). Accessed July 23, 2016.
  27. Thackray H, Tiffit C. Fetal alcohol syndrome. *Pediatr Rev*. 2001;22(2):47-55.
  28. Adams DJ, Clark DA. Common genetic and epigenetic syndromes. *Pediatr Clin North Am*. 2015;62(2):411-426.
  29. Streissguth AP, Bookstein FL, Barr HM, Sampson PD, O'Malley K, Young JK. Risk factors for adverse life outcomes in fetal alcohol syndrome and fetal alcohol effects. *J Dev Behav Pediatr*. 2004;25(4):228-238.
  30. Pruett D, Waterman EH, Caughey AB. Fetal alcohol exposure: consequences, diagnosis, and treatment. *Obstet Gynecol Surv*. 2013;68(1):62-69.
  31. Carter RC, Jacobson SW, Moltano CD, Chiodo LM, Viljoen D, Jacobson JL. Effects of prenatal alcohol exposure on infant visual acuity. *J Pediatr*. 2005;147(4):473-479.
  32. Fuglestad AJ, Fink BA, Eckerle JK, et al. Inadequate intake of nutrients essential for neurodevelopment in children with fetal alcohol spectrum disorders (FASD). *Neurotoxicol Teratol*. 2013;39:128-132.
  33. Young JK, Giesbrecht HE, Eskin MN, Aliani M, Suh M. Nutrition implications for fetal alcohol spectrum disorder. *Adv Nutr*. 2014;5(6):675-692.
  34. Burd L, Cotsonas-Hassler TM, Martsolf JT, Kerbeshian J. Recognition and management of fetal alcohol syndrome. *Neurotoxicol Teratol*. 2003;25(6):681-688.
  35. Fryer SL, McGee CL, Matt GE, Riley EP, Mattson SN. Evaluation of psychological conditions in children with heavy prenatal alcohol exposure. *Pediatrics*. 2007;119(3):e733-e741.
  36. O'Connor MJ, Shah B, Whaley S, Cronin P, Gunderson B, Graham J. Psychiatric illness in a clinical sample of children with prenatal alcohol exposure. *Am J Drug Alcohol Abuse*. 2002;28(4):743-754.
  37. Davis K, Desrocher M, Moore T. Fetal alcohol spectrum disorder: a review of neurodevelopmental findings and interventions. *J Dev Phys Disabil*. 2011;23(2):143-167.
  38. Oesterheld JR, Kofoed L, Tervo R, Fogas B, Wilson A, Fiechtner H. Effectiveness of methylphenidate in Native American children with fetal alcohol syndrome and attention deficit/hyperactivity disorder: a controlled pilot study. *J Child Adolesc Psychopharmacol*. 1998;8(1):39-48.
  39. Freunsch I, Feldmann R. Young adults with fetal alcohol syndrome (FAS): social, emotional and occupational development. *Klin Padiatr*. 2011;223(1):33-37.
  40. Rangmar J, Hjern A, Vinnerljung B, Strömland K, Aronson M, Fahlke C. Psychosocial outcomes of fetal alcohol syndrome in adulthood. *Pediatrics*. 2015;135(1):e52-e58.
  41. Williams JF, Smith VC; Committee on Substance Abuse. Fetal alcohol spectrum disorders. *Pediatrics*. 2015;136(5):e1395-e1406.
  42. American College of Obstetricians and Gynecologists; Committee on Health Care for Underserved Women. Committee opinion no. 496: at-risk drinking and alcohol dependence: obstetric and gynecologic implications. *Obstet Gynecol*. 2011;118(2 pt 1):383-388.
  43. American Academy of Family Physicians. Substance abuse and addiction. <http://www.aafp.org/about/policies/all/substance-abuse.html>. Accessed December 9, 2016.
  44. Green P, McKnight-Eily L, Tan C, Mejia R, Denny C. Vital signs: alcohol-exposed pregnancies—United States, 2011-2013. *MMWR Morb Mortal Wkly Rep*. 2016;65(4):91-97.
  45. Carson G, Cox LV, Crane J, et al.; Society of Obstetricians and Gynaecologists of Canada. Alcohol use and pregnancy consensus clinical guidelines. *J Obstet Gynaecol Can*. 2010;32(8 suppl 3):S1-S31.
  46. Chiodo LM, Delaney-Black V, Sokol RJ, Janisse J, Pardo Y, Hannigan JH. Increased cut-point of the TACER-3 screen reduces false positives without losing sensitivity in predicting risk alcohol drinking in pregnancy. *Alcohol Clin Exp Res*. 2014;38(5):1401-1408.
  47. Committee opinion no. 633: alcohol abuse and other substance use disorders: ethical issues in obstetric and gynecologic practice. *Obstet Gynecol*. 2015;125(6):1529-1537.
  48. Chang G, McNamara TK, Orav EJ, et al. Brief intervention for prenatal alcohol use: a randomized trial. *Obstet Gynecol*. 2005;105(5 pt 1):991-998.
  49. O'Connor MJ, Whaley SE. Brief intervention for alcohol use by pregnant women. *Am J Public Health*. 2007;97(2):252-258.



### eTable A. Diagnosis of Alcohol-Related Birth Defects

---

Documented prenatal alcohol exposure

At least 1 of the following specific major malformations known to be the result of prenatal alcohol exposure:

Auditory: conductive and/or neurosensory hearing loss

Cardiac: aberrant great vessels, atrial septal defect, conotruncal heart defects, ventricular septal defect

Musculoskeletal: flexion contractures, radioulnar synostosis, scoliosis, vertebral segmentation defects

Ophthalmologic: optic nerve hypoplasia, ptosis, retinal vascular anomalies, strabismus

Renal: aplastic/dysplastic/hypoplastic kidneys, horseshoe kidney, ureteral duplications

---

*Information from Hoyme HE, Kalberg WO, Elliott AJ, et al. Updated clinical guidelines for diagnosing fetal alcohol spectrum disorders. Pediatrics. 2016;138(2):e20154256.*