



Fanconi Anemia

Synonym: Fanconi Pancytopenia

Parinda A Mehta, MD¹ and Jakub Tolar, MD, PhD²

Created: February 14, 2002; Revised: March 8, 2018.

Summary

Clinical characteristics

Fanconi anemia (FA) is characterized by physical abnormalities, bone marrow failure, and increased risk for malignancy. Physical abnormalities, present in approximately 75% of affected individuals, include one or more of the following: short stature, abnormal skin pigmentation, skeletal malformations of the upper and lower limbs, microcephaly, and ophthalmic and genitourinary tract anomalies. Progressive bone marrow failure with pancytopenia typically presents in the first decade, often initially with thrombocytopenia or leukopenia. The incidence of acute myeloid leukemia is 13% by age 50 years. Solid tumors – particularly of the head and neck, skin, gastrointestinal tract, and genitourinary tract – are more common in individuals with FA.

Diagnosis/testing

The diagnosis of FA is established in a proband with increased chromosome breakage and radial forms on cytogenetic testing of lymphocytes with diepoxybutane (DEB) and mitomycin C (MMC). The diagnosis is confirmed by identification of one of the following:

- Biallelic pathogenic variants in one of the 19 genes known to cause autosomal recessive FA
- A heterozygous pathogenic variant in *RAD51*, known to cause autosomal dominant FA
- A hemizygous pathogenic variant in *FANCB*, known to cause X-linked FA

Management

Treatment of manifestations: Administration of oral androgens (e.g., oxymetholone) improves blood counts (red cell and platelets) in approximately 50% of individuals with FA; administration of G-CSF improves the neutrophil count in some; hematopoietic stem cell transplantation (HSCT) is the only curative therapy for the hematologic manifestations of FA, but the high risk for solid tumors remains and may even be increased in those undergoing HSCT. All these treatments have potential significant toxicity. Early detection and surgical removal remains the mainstay of therapy for solid tumors.

Author Affiliations: 1 Professor, Division of Bone Marrow Transplantation and Immune Deficiency, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio; Email: parinda.mehta@cchmc.org. 2 Professor, Division of Blood and Marrow Transplantation, Stem Cell Institute, University of Minnesota, Minneapolis, Minnesota.

Prevention of primary manifestations: Human papilloma virus (HPV) vaccination to reduce the risk of gynecologic cancer in females, and possibly reduce the risk of oral cancer in all individuals.

Prevention of secondary complications: T-cell depletion of the donor graft to minimize the risk of graft vs host disease; conditioning regimen without radiation prior to HSCT to reduce the risk of subsequent solid tumors.

Surveillance: Annual evaluation with a multidisciplinary team including an endocrinologist; monitoring for evidence of bone marrow failure (regular blood counts; at least annual bone marrow aspirate/biopsy to evaluate morphology, cellularity, and cytogenetics); for those receiving androgen therapy, monitoring liver function tests and regular ultrasound examination of the liver; monitoring for solid tumors (oropharyngeal and gynecologic examinations).

Agents/circumstances to avoid: Transfusions of red cells or platelets for persons who are candidates for HSCT; family members as blood donors if HSCT is being considered; blood products that are not filtered (leukodepleted) or irradiated; toxic agents that have been implicated in tumorigenesis; unsafe sex practices, which increase the risk of HPV-associated malignancy; radiographic studies solely for the purpose of surveillance (i.e., in the absence of clinical indications).

Evaluation of relatives at risk: DEB/MMC testing or molecular genetic testing (if the family-specific pathogenic variants are known) of all sibs of a proband for early diagnosis, treatment, and monitoring for physical abnormalities, bone marrow failure, and related cancers.

Genetic counseling

Fanconi anemia (FA) can be inherited in an autosomal recessive manner, an autosomal dominant manner (*RAD51*-related FA), or an X-linked manner (*FANCB*-related FA).

Autosomal recessive FA: Each sib of an affected individual has a 25% chance of inheriting both pathogenic variants and being affected, a 50% chance of inheriting one pathogenic variant and being a carrier, and a 25% chance of inheriting both normal alleles and not being a carrier. Carriers (heterozygotes) for autosomal recessive FA are asymptomatic.

Autosomal dominant FA: Given that all affected individuals with *RAD51*-related FA reported to date have the disorder as a result of a *de novo* *RAD51* pathogenic variant, the risk to other family members is presumed to be low.

X-linked FA: For carrier females the chance of transmitting the pathogenic variant in each pregnancy is 50%; males who inherit the pathogenic variant will be affected; females who inherit the pathogenic variant will be carriers and will usually not be affected.

Carrier testing for at-risk relatives (for autosomal recessive and X-linked FA) and prenatal testing for pregnancies at increased risk are possible if the pathogenic variant(s) in the family are known.

Diagnosis

Recommendations for diagnosis were agreed upon at a 2013 consensus conference (see [Fanconi Anemia: Guidelines for Diagnosis and Management](#)).

Suggestive Findings

Fanconi anemia (FA) **should be suspected** in individuals with the following clinical and laboratory features.

Physical features (in ~75% of affected persons)

- Prenatal and/or postnatal short stature

- Abnormal skin pigmentation (e.g., café au lait macules, hypopigmentation)
- Skeletal malformations (e.g., hypoplastic thumb, hypoplastic radius)
- Microcephaly
- Ophthalmic anomalies
- Genitourinary tract anomalies

Laboratory findings

- Macrocytosis
- Increased fetal hemoglobin (often precedes anemia)
- Cytopenia (especially thrombocytopenia, leukopenia and neutropenia)

Pathology findings

- Progressive bone marrow failure
- Adult-onset aplastic anemia
- Myelodysplastic syndrome (MDS)
- Acute myelogenous leukemia (AML)
- Early-onset solid tumors (e.g., squamous cell carcinomas of the head and neck, esophagus, and vulva; cervical cancer; and liver tumors)
- Inordinate toxicities from chemotherapy or radiation

Establishing the Diagnosis

The diagnosis of FA is **established** in a proband with the following findings:

- Increased chromosome breakage and radial forms on cytogenetic testing of lymphocytes with diepoxybutane (DEB) and mitomycin C (MMC)

Note: (1) The background rate of chromosome breakage in control chromosomes is more variable with MMC; thus, some centers use DEB while other centers use both DEB and MMC. (2) If results of lymphocyte testing are normal or inconclusive and mosaicism is suspected, testing can be performed on an alternative cell type, such as skin fibroblasts. See [Fanconi Anemia: Guidelines for Diagnosis and Management](#).

- Identification of biallelic pathogenic variants in one of the 18 genes known to cause autosomal recessive FA, or a heterozygous pathogenic variant in *RAD51* known to cause autosomal dominant FA, or a hemizygous pathogenic variant in *FANCB*, known to cause X-linked FA (see Table 1)

Molecular testing approaches can include **single-gene testing**, use of a **multigene panel**, and **more comprehensive genomic testing**:

- **Single-gene testing.** Sequence analysis of *FANCA* can be performed first, followed by gene-targeted *FANCA* deletion/duplication analysis if only one or no pathogenic variant is found.
- **A multigene panel** that includes the genes in Table 1a and Table 1b and other genes of interest (see Differential Diagnosis) may be considered next if single-gene testing does not identify a *FANCA* pathogenic variant. Note: (1) The genes included in the panel and the diagnostic sensitivity of the testing used for each gene vary by laboratory and are likely to change over time. (2) Some multigene panels may include genes not associated with the condition discussed in this *GeneReview*; thus, clinicians need to determine which multigene panel provides the best opportunity to identify the genetic cause of the condition at the most reasonable cost while limiting identification of pathogenic variants in genes that do

not explain the underlying phenotype. (3) Methods used in a panel may include sequence analysis, deletion/duplication analysis, and/or other non-sequencing-based tests.

For an introduction to multigene panels click [here](#). More detailed information for clinicians ordering genetic tests can be found [here](#).

- **More comprehensive genomic testing** (when available) including exome sequencing and genome sequencing may be considered if serial single-gene testing (and/or use of a multigene panel) fails to confirm a diagnosis in an individual with features of Fanconi anemia. Such testing may provide or suggest a diagnosis not previously considered (e.g., mutation of a different gene or genes that results in a similar clinical presentation).

For an introduction to comprehensive genomic testing click [here](#). More detailed information for clinicians ordering genomic testing can be found [here](#).

Table 1a. Molecular Genetics of Fanconi Anemia: Most Common Genetic Causes

Gene ^{1, 2}	Complementation Group ³	% of FA Attributed to Pathogenic Variants in This Gene ⁴	Proportion of Pathogenic Variants ⁵ Detected by Method	
			Sequence analysis ⁶	Gene-targeted deletion/duplication analysis ⁷
<i>BRCA2</i>	FA-D1	~3%	>90%	None reported
<i>BRIP1</i>	FA-J	~2%	>90%	None reported
<i>FANCA</i>	FA-A	60%-70%	~60%	>20 deletion/duplication variants reported; see HGMD
<i>FANCB</i>	FA-B	~2%	Unknown	Deletion/duplication variants reported; see HGMD
<i>FANCC</i>	FA-C	~14%	Unknown	Several deletion/duplication variants reported; see HGMD
<i>FANCD2</i>	FA-D2	~3%	Unknown	Multiple deletion/duplication variants reported; see HGMD
<i>FANCE</i>	FA-E	~3%	>90%	None reported
<i>FANCF</i>	FA-F	~2%	Unknown	Deletion/duplication variants reported; see HGMD
<i>FANCG</i>	FA-G	~10%	>90%	None reported

Table 1a. continued from previous page.

Gene ^{1, 2}	Complementation Group ³	% of FA Attributed to Pathogenic Variants in This Gene ⁴	Proportion of Pathogenic Variants ⁵ Detected by Method	
			Sequence analysis ⁶	Gene-targeted deletion/duplication analysis ⁷
<i>FANCI</i>	FA-I	~1%	Unknown	Deletion/duplication variant reported; see HGMD

Pathogenic variants of any one of the genes included in this table account for >1% of FA

1. Genes are listed in alphabetic order.

2. See Table A. Genes and Databases for chromosome locus and protein.

3. Prior to identification of the genes, complementation groups were defined based on somatic cell-based methods. While complementation analysis testing has been supplanted by multigene panels; this terminology continues to be used in some contexts.

4. Shimamura & Alter [2010]

5. See Molecular Genetics for information on pathogenic variants detected.

6. Sequence analysis detects variants that are benign, likely benign, of uncertain significance, likely pathogenic, or pathogenic. Pathogenic variants may include small intragenic deletions/insertions and missense, nonsense, and splice site variants; typically, exon or whole-gene deletions/duplications are not detected. For issues to consider in interpretation of sequence analysis results, click [here](#).

7. Gene-targeted deletion/duplication analysis detects intragenic deletions or duplications. Methods that may be used include: quantitative PCR, long-range PCR, multiplex ligation-dependent probe amplification (MLPA), and a gene-targeted microarray designed to detect single-exon deletions or duplications.

Table 1b. Molecular Genetics of Fanconi Anemia: Less Common Genetic Causes

Gene ^{1, 2, 3}	Complementation Group ⁴	Comment
<i>ERCC4</i>	FA-Q	2 individuals w/FA & biallelic <i>ERCC4</i> pathogenic variants reported [Bogliolo et al 2013]; functional studies of <i>ERCC4</i> missense variants reported by Osorio et al [2013]
<i>FANCL</i>	FA-L	13 <i>FANCL</i> pathogenic variants reported [Chandrasekharappa et al 2013, Nicchia et al 2015, Vetro et al 2015]
<i>FANCM</i>	FA-M	Assignment of a formal complementation group for persons w/ <i>FANCM</i> pathogenic variants still controversial as only 1 reference family/cell line has been identified & that cell line has been determined to have biallelic pathogenic variants in both <i>FANCA</i> & <i>FANCM</i> . Of note, under experimental conditions specific knockdown of <i>FANCM</i> alone results in an FA phenotype [Singh et al 2009].
<i>MAD2L2</i>	FA-V	1 individual w/homozygous pathogenic variants reported [Bluteau et al 2016]
<i>PALB2</i>	FA-N	14 <i>PALB2</i> pathogenic variants reported incl a deletion of exons 1-10 [Reid et al 2007, Xia et al 2007, Byrd et al 2016]
<i>RAD51</i>	FA-R	2 individuals w/features of FA & a <i>de novo</i> <i>RAD51</i> pathogenic variant reported [Ameziane et al 2015, Wang et al 2015]
<i>RAD51C</i>	FA-O	Assignment of a formal complementation group for persons w/ <i>RAD51C</i> pathogenic variants still controversial as only 1 reference consanguineous family identified [Vaz et al 2010]
<i>RFWD3</i>	FA-W	1 individual w/features of FA & compound heterozygous pathogenic variants in <i>RFWD3</i> reported [Knies et al 2017]
<i>SLX4</i>	FA-P	Assignment of a formal complementation group for persons w/ <i>SLX4</i> pathogenic variants still controversial as only a handful of reference families have been identified & <i>SLX4</i> biology falls outside previously characterized FA proteins [Kim et al 2011, Stoepker et al 2011]
<i>UBE2T</i>	FA-T	1 individual w/biallelic <i>UBE2T</i> pathogenic variants incl a large paternal deletion & a maternal duplication reported [Rickman et al 2015, Virts et al 2015]

Table 1b. continued from previous page.

Gene ^{1, 2, 3}	Complementation Group ⁴	Comment
<i>XRCC2</i>	FA-U	1 individual w/homozygous pathogenic variants reported [Park et al 2016]

Pathogenic variants of any one of the genes listed in this table are reported in only a few families (i.e., <1% of FA)

1. Genes are listed in alphabetic order.

2. See Table A. Genes and Databases for chromosome locus and protein.

3. Genes are not described in detail in Molecular Genetics but may be included [here](#) (pdf).

4. Prior to identification of the genes, complementation groups were defined based on somatic cell-based methods this terminology continues to be used in some contexts.

Clinical Characteristics

Clinical Description

The primary clinical features of Fanconi anemia (FA) include physical features, progressive bone marrow failure manifest as pancytopenia, and cancer susceptibility; however, some individuals with FA have neither physical abnormalities nor bone marrow failure.

Physical features occur in approximately 75% of individuals with FA.

- Growth deficiency: prenatal and/or postnatal short stature, low birth weight
- Abnormal skin pigmentation (40%): generalized hyperpigmentation; café au lait macules, hypopigmentation
- Skeletal malformations of upper limbs, unilateral or bilateral (35%):
 - Thumbs (35%): absent, hypoplastic, bifid, duplicated, triphalangeal, long, proximally placed
 - Radii (7%): absent or hypoplastic (only with abnormal thumbs), absent or weak pulse
 - Hands (5%): flat thenar eminence, absent first metacarpal, clinodactyly, polydactyly
 - Ulnae (1%): dysplastic, short
- Skeletal malformations of lower limbs (5%)
 - Syndactyly, abnormal toes, club feet
 - Congenital hip dislocation
- Microcephaly (20%)
- Ophthalmic (20%): microphthalmia, cataracts, astigmatism, strabismus, epicanthal folds, hypotelorism, hypertelorism, ptosis
- Genitourinary tract anomalies:
 - Renal (20%): horseshoe, ectopic, pelvic, hypoplastic, dysplastic, or absent kidney; hydronephrosis or hydroureter
 - Males (25%). Hypospadias, micropenis, cryptorchidism, anorchia, hypo- or azoospermia, reduced fertility
 - Females (2%). bicornuate or uterus malposition, small ovaries

Note: Pregnancy is possible in females, whether or not they have undergone hematopoietic stem cell transplantation.
- Endocrine: hypothyroidism, glucose/insulin abnormalities

- Hearing loss, usually conductive secondary to middle ear bony anomalies; abnormal ear shape: dysplastic, narrow ear canal, abnormal pinna (10%)
- Congenital heart defect (6%): patent ductus arteriosus, atrial septal defect, ventricular septal defect, coarctation of the aorta, truncus arteriosus, situs inversus
- Gastrointestinal (5%): esophageal, duodenal, or jejunal atresia, imperforate anus, tracheoesophageal fistula, annular pancreas, malrotation
- Central nervous system (3%): small pituitary, pituitary stalk interruption syndrome, absent corpus callosum, cerebellar hypoplasia, hydrocephalus, dilated ventricles
- Other
 - Facial features (2%): triangular, micrognathia, mid-face hypoplasia
 - Spine anomalies (2%): spina bifida, scoliosis, hemivertebrae, rib anomalies, coccygeal aplasia
 - Neck anomalies (1%): Sprengel deformity, Klippel-Feil anomaly, short or webbed neck, low hairline

Note: Percentages are calculated from 2,000 individuals reported in the literature from 1927 to 2009. Frequencies are approximate, since many reports did not mention physical descriptions.

Developmental delay and/or intellectual disability (10%)

Bone marrow failure. The age of onset is highly variable, even among sibs. An analysis of 754 individuals with pathogenic variants in *FANCA*, *FANCC*, and *FANCG* identified an average age of onset of 7.6 years. Rarely, bone marrow failure can present in infants and small children [Shimamura & Alter 2010]. The risk of developing any hematologic abnormality is 90% by age 40 years [Kutler et al 2003].

- Thrombocytopenia or leukopenia usually precede anemia. These are commonly associated with macrocytosis and elevated fetal hemoglobin.
- Pancytopenia generally worsens over time.
- Sweet syndrome (neutrophilic skin infiltration) was associated with progression of hematologic disease in six out of seven individuals with FA [Giulino et al 2011].
- The severity of bone marrow failure can be classified by the degree of cytopenia(s) (Table 3). Importantly, to meet these criteria for marrow failure, the cytopenias must be persistent and unexplained by other causes.

Table 2. Severity of Bone Marrow Failure in Fanconi Anemia

	Mild	Moderate	Severe
Absolute neutrophil count (ANC)	<1,500/mm ³	<1,000/mm ³	<500/mm ³
Platelet count	150,000-50,000/mm ³	<50,000/mm ³	<30,000/mm ³
Hemoglobin (Hb) level	≥8 g/dL	<8 g/dL	<8 g/dL

Cancer susceptibility. The relative risk for acute myelogenous leukemia (AML) is increased approximately 500-fold [Rosenberg et al 2008, Alter et al 2010, Tamary et al 2010]. In a competing risk analysis of the combined cohorts, the cumulative incidence of AML was 13% by age 50 years, with most individuals diagnosed between ages 15 and 35 years.

An increased risk of developing myelodysplastic syndrome (MDS)/AML is associated with monosomy 7 and most 7q deletions. Clonal amplifications of chromosome 3q26-q29 were reported in association with an increased risk of progression to MDS/AML [Neitzel et al 2007, Mehta et al 2010].

Solid tumors may be the first manifestation of FA in individuals who have no birth defects and have not experienced bone marrow failure.

- Head and neck squamous cell carcinomas (HNSCCs) are the most common solid tumor in individuals with FA. The incidence is 500- to 700-fold higher than in the general population. The HNSCCs in FA show distinct differences compared to HNSCCs seen in the general population. HNSCCs:
 - Occur at an earlier age (20-40 years) than in the general population;
 - Are most commonly in the oral cavity (e.g., tongue);
 - Present at an advanced stage;
 - Respond poorly to therapy.
- Individuals with FA are at increased risk for second primary cancers in the skin and genitourinary tract. The pattern of second primaries resembles that observed in HPV-associated HNSCC in the general population [Morris et al 2011].
- Individuals with FA receiving androgen treatment for bone marrow failure are also at increased risk for liver tumors.

Phenotype Correlations by Gene

BRCA2. Biallelic pathogenic variants in *BRCA2* are associated with early-onset acute leukemia and solid tumors [Hirsch et al 2004, Wagner et al 2004, Myers et al 2012]. The cumulative probability of any malignancy was 97% by age six years, including AML, medulloblastoma, and Wilms tumor [Alter et al 2007].

FANCG. Pathogenic variants in *FANCG* may be associated with severe marrow failure and a higher incidence of leukemia compared to *FANCC* [Faivre et al 2000].

PALB2. Solid tumors (e.g., medulloblastoma, Wilms tumor) are associated with *PALB2* pathogenic variants [Reid et al 2007].

Genotype-Phenotype Correlations

The clinical spectrum of FA remains heterogenous. There are no clearcut genotype-phenotype correlations [Neveling et al 2009]. In general, null variants lead to a more severe phenotype (e.g., congenital anomalies, early-onset bone marrow failure, and MDS/AML) than hypomorphic variants.

BRCA2. All persons with an IVS7 pathogenic variant in *BRCA2* developed AML by age three years; those with other *BRCA2* pathogenic variants who developed AML did so by age six years [Alter 2006].

FANCA. Individuals who are homozygous for null pathogenic variants in *FANCA* may have earlier onset of anemia and higher incidence of leukemia than individuals with pathogenic variants that permit production of an abnormal *FANCA* protein [Faivre et al 2000].

FANCC

- c.456+4A>T, p.Arg548Ter, and p.Leu554Pro are associated with earlier onset of hematologic abnormalities and more severe congenital anomalies than other pathogenic variants, such as del22G [Faivre et al 2005].
- p.Asp23IlefsTer23 and p.Gln13Ter are associated with a lower risk for congenital anomalies and later progression to bone marrow failure [Yamashita et al 1996, Gillio et al 1997].
- c.456+4A>T results in a milder phenotype in Japanese individuals than in Ashkenazi Jewish individuals [Futaki et al 2000].

Prevalence

Fanconi anemia (FA) is the most common genetic cause of aplastic anemia and one of the most common genetic causes of hematologic malignancy.

The ratio of males to females is 1.2:1 (p<0.001 vs expected 1.00).

Rosenberg et al [2011] showed higher carrier rates for FA than previously reported. Carrier frequency was 1:181 in North Americans and 1:93 in Israel. Specific populations have founder variants with increased carrier frequencies (<1:100), including Ashkenazi Jews (*FANCC*, *BRCA2*), northern Europeans (*FANCC*), Afrikaners (*FANCA*), sub-Saharan Blacks (*FANCG*), Spanish Gypsies (*FANCA*), and others.

Genetically Related (Allelic) Disorders

Hereditary breast and ovarian cancer is associated with heterozygous pathogenic variants in *BRCA2*.

Heterozygous pathogenic variants in *BRIP1* and *PALB2* have been implicated in breast cancer and ovarian cancer predisposition [Seal et al 2006, Rahman et al 2007, Rafnar et al 2011].

Heterozygous pathogenic variants in *PALB2* have been implicated in pancreatic cancer predisposition [Jones et al 2009].

Biallelic pathogenic variants in *ERCC4* are associated with [xeroderma pigmentosum](#), [Cockayne syndrome](#), and XFE progeroid syndrome.

Differential Diagnosis

Cells derived from individuals with other chromosome breakage syndromes, such as [Bloom syndrome](#) or [ataxia-telangiectasia](#), may also exhibit high rates of spontaneous chromosome breakage; however, only FA cells exhibit increased chromosome breakage in response to diepoxybutane (DEB).

[Nijmegen breakage syndrome](#) (NBS), characterized by short stature, progressive microcephaly with loss of cognitive skills, premature ovarian failure in females, recurrent sinopulmonary infections, and an increased risk for cancer (particularly lymphoma), may also manifest increased chromosome breakage with mitomycin C (MMC) [Chrzanowska et al 2012]. Inheritance is autosomal recessive. NBS may be distinguished from FA by *NBN* molecular genetic testing, which identifies pathogenic variants in almost 100% of individuals with NBS.

Seckel syndrome (OMIM [PS210600](#)), characterized by growth retardation, microcephaly with intellectual disability, and a characteristic "bird-headed" facial appearance, may also show increased chromosome breakage with DNA cross-linking agents (MMC, DEB) [Andreassen et al 2004]. Some individuals with Seckel syndrome also develop pancytopenia and/or AML. Inheritance is autosomal recessive. Biallelic pathogenic variants in *ATR*, *NIN*, *ATRIP*, *RBBP8*, *CEP152*, *CENPJ*, and *CEP63* are causative.

Other disorders including [neurofibromatosis 1](#) (which could be considered because of café au lait macules), [TAR syndrome](#) (thrombocytopenia with absent radii), and VACTERL association (radial ray defects) (OMIM [192350](#)) can be distinguished from FA by testing for chromosome breakage with DEB and MMC.

Management

Evaluations Following Initial Diagnosis

To establish the extent of disease and management requirements in an individual diagnosed with Fanconi anemia (FA), the following evaluations are recommended:

- Evaluation by a hematologist, to include complete blood count, fetal hemoglobin, full blood typing, blood chemistries (assessing liver, kidney, and iron status), and bone marrow aspirate for cell morphology, FISH and cytogenetics, as well as biopsy for cellularity

Note: The bone marrow of individuals with FA can exhibit signs of dysplasia, such as nuclear/cytoplasmic dys-synchrony, hypo-lobulated megakaryocytes, and bi-nucleated erythroid cells. These features must be

distinguished from true forms of MDS by a hematopathologist experienced in the evaluation of MDS in individuals with FA.

- HLA typing of the affected individual, sibs, and parents for consideration of hematopoietic stem cell transplantation
- Examination by an ophthalmologist
- Ultrasound examination of the kidneys and urinary tract
- Formal hearing evaluation
- Echocardiogram
- Referral to an endocrinologist
- Developmental assessment (particularly important for toddlers and school-age children)
- Referrals as indicated to an otolaryngologist, hand surgeon, gastroenterologist, gynecologist, and urologist
- Evaluation by a clinical geneticist and genetic counseling

Treatment of Manifestations

Recommendations for treatment were agreed upon at a 2014 consensus conference ([full text](#)).

Androgens improve (at least transiently) the red cell and platelet counts in approximately 50% of individuals. Androgen therapy can be considered when the hemoglobin drops below 8 g/dL or the platelet count falls below 30,000/mm³ ("severe" – see Table 2). Although only 10%-20% of individuals receiving continuous low-dose androgen therapy are long-term responders, this option can be particularly useful for individuals who do not have access to or are not ready for hematopoietic stem cell transplant (HSCT), or to individuals for whom a suitable donor is not available.

- Oxymetholone, given orally at a starting dose of 2 mg/kg/day, may be increased up to 5 mg/kg/day.
- Doses may be slowly tapered to the minimal effective dose with careful monitoring of the blood counts.
- Other synthetic androgens used in FA include stanzolol in Asia, and oxandrolone and danazol in North America.

Side effects of androgen administration include virilization and liver toxicity such as elevated liver enzymes, cholestasis, peliosis hepatis (vascular lesion with multiple blood-filled cysts), and hepatic tumors. Individuals taking androgens should be monitored for liver tumors and undergo regular liver function tests (LFT) for abnormalities. Blood tests for LFTs should be performed every three to six months; liver ultrasound should be performed every six to 12 months. If no response is seen after three to four months, androgens should be discontinued [Scheckenbach et al 2012, Rose et al 2014].

Granulocyte colony-stimulating factor (G-CSF) improves the neutrophil count in some individuals. G-CSF dose should be titrated to the lowest possible dose and frequency to keep ANC above 1,000/mm³. Note: (1) A bone marrow aspirate and biopsy should be performed prior to the initiation of G-CSF and monitored every six months throughout treatment, given the theoretic risk of stimulating the growth of a leukemic clone. (2) G-CSF should be administered in consultation with an FA expert.

Hematopoietic stem cell transplantation (HSCT) is the only curative therapy for the hematologic manifestations, including aplastic anemia, myelodysplastic syndrome, and acute leukemia. Ideally, HSCT is performed prior to onset of MDS/AML and before multiple transfusions [MacMillan & Wagner 2010, Mehta et

al 2010]. Individuals with FA are sensitive to chemotherapy and radiation, need special transplant regimens, and should be cared for and transplanted at centers with the most experience in HSCT in FA.

A multi-institutional study reported a one-year probability of overall survival of 80% in 45 individuals with FA transplanted for marrow failure and/or MDS, using alternative donors (including mismatched related and unrelated donors) and chemotherapy-only preparative regimen. Survival for individuals younger than age ten years transplanted for marrow failure was even better, at 91.3% ($\pm 5.9\%$) [Mehta et al 2017].

Fludarabine reduced the incidence of graft failure and allowed for removal of radiation from the preparative regimens in a matched sib donor setting [MacMillan et al 2015].

MDS/AML treatment remains challenging. Options include chemotherapy, HSCT with or without prior induction chemotherapy, and investigational trials. Chemotherapy should be undertaken in coordination with centers experienced with FA, as it can cause severe, prolonged, or irreversible myelosuppression. Plans for HSCT should be in place prior to starting chemotherapy. Published reports of chemotherapy regimens for AML in individuals with FA are sparse and limited by the unclear benefit to the overall outcome due to the lack of longitudinal follow up [Mehta et al 2007, Talbot et al 2014, Beier et al 2015].

Solid tumors. Prompt, aggressive workup for any symptoms suggestive of a malignancy is indicated. Early detection and surgical removal remains the mainstay of therapy. Treatment is challenging secondary to the increased toxicity associated with chemotherapy and radiation in FA. Data is limited on use of chemotherapy at standard doses or reduced doses and schedules in individuals with FA, and there are reports of severe or fatal toxicities and poor treatment outcomes [Masserot et al 2008, Hosoya et al 2010, Tan et al 2011, Spanier et al 2012]. Individuals diagnosed with a genital tract cancer should be referred to a gynecologic oncologist immediately, and care should be coordinated with FA experts.

Prevention of Primary Manifestations

Human papilloma virus (HPV) vaccination should be initiated at age nine years in order to reduce the risk of gynecologic cancer in females, and possibly reduce the risk of oral cancer in all individuals.

Prevention of Secondary Complications

Individuals with FA treated with HSCT who developed graft vs host disease (GVHD) had a 28% incidence of head and neck cancers in the ten years following treatment (vs 0% in those without GVHD); this finding points to the importance of minimizing the risk of GVHD [Guardiola et al 2004]. Increased risk for GVHD observed in earlier studies was reduced significantly by T-cell depletion of the donor graft [Chaudhury et al 2008, MacMillan et al 2015].

Individuals successfully treated with HSCT are at increased risk for solid tumors, in addition to the baseline increased risk [Rosenberg et al 2005]. Due to the known contribution of radiation to the long-term complication of secondary solid tumors most recent efforts have focused on using a conditioning regimen without radiation even in an unrelated donor setting. German, Brazilian, and US groups now report excellent outcomes with alternative donors with a "chemotherapy-only" preparative regimen in single-center studies. The study from Germany showed 88% survival and normal hematopoiesis at a median follow up of 30 months [Bonfim et al 2015, Chao et al 2015]. A prospective multi-institutional US study also showed similar excellent outcomes. One-year probabilities of overall and disease-free survival for the entire cohort, including patients with myeloid malignancy and those receiving mismatched related/haploidentical grafts, were 80% and 77.7% respectively at a median follow-up of 41 months. All young children (age <10 years) undergoing HSCT for marrow failure using low-dose busulfan-containing regimen survived [Mehta et al 2017].

Surveillance

See 2014 consensus guidelines [Frohnmayr et al 2014] ([full text](#)).

- Annual evaluation with a multidisciplinary team including an endocrinologist
- Regular blood counts, every three to four months while stable and more often as needed
- Bone marrow aspirate/biopsy at least annually to evaluate morphology, cellularity (from the biopsy), FISH, and cytogenetics (the latter two for emergence of a malignant clone). Individuals on GCSF need to have a bone marrow aspirate/biopsy every six months, if possible.
- In individuals who develop Sweet syndrome (neutrophilic skin infiltration), prompt investigation for hematologic disease progression including bone marrow evaluation

Notes: (1) Progressively changing blood counts without a potential cause (e.g., acute infection or suppression from medication) require immediate evaluation with a complete blood count and bone marrow examination with FISH and cytogenetics. (2) It is important to recognize that rising blood counts can be due to either the development of MDS/AML or, rarely, reversion of a germline mutation in a stem cell, which repopulates the marrow with normal cells (somatic stem cell mosaicism). These individuals may require immediate HSCT (for MDS/AML) or continued close monitoring with complete blood counts at least every one to two months and a bone marrow examination with cytogenetics every six months.

Individuals receiving androgen therapy

- Liver function tests every three to six months
- Liver ultrasound examination every six to 12 months for androgen-related changes, including tumors

Cancer surveillance

- Annual gynecologic assessment for genital lesions beginning at age 13. Thorough vulvo-vaginal examinations and Pap smear can begin when women become sexually active or by age 18 years, whichever is earlier. Suspicious genital tract lesions should be biopsied.
- Examination every six months for oral, head, and neck cancers beginning by age nine to ten years. Screening should be performed by a dentist, oral surgeon, or ENT familiar with FA. Nasolaryngoscopy starting at age ten years, or within the first year after HSCT. Individuals with difficulty or pain with swallowing should be evaluated for esophageal cancer.
- For individuals with a history of premalignant or malignant lesions: surveillance examinations every two to three months
- For individuals with biallelic pathogenic variants in *BRCA2*: screening for neuroblastomas, brain tumors, and kidney tumors every six months (see also Autosomal Recessive FA, Risk to Family Members)

Agents/Circumstances to Avoid

Blood transfusions. Blood products should be cytomegalovirus (CMV)-safe and irradiated. To reduce the chances of sensitization, family members must not act as blood donors. Once an individual requires transfusions, he/she should be referred for transplantation.

Toxic agents to avoid include smoking, second-hand smoke, and alcohol, which have been implicated in tumorigenesis.

Unsafe sex practices increase the risk for HPV-associated malignancy.

Radiographic studies for the purpose of surveillance should be minimized in the absence of clinical indications. However, baseline skeletal surveys may be considered, in order to document bony anomalies that may lead to problems with age, such as anomalies of the wrist, hip, and vertebrae.

Evaluation of Relatives at Risk

It is appropriate to evaluate all sibs of an affected individual in order to identify as early as possible those who would benefit from appropriate monitoring for physical abnormalities, bone marrow failure, and related cancers.

- DEB/MMC testing can be used to clarify the disease status of at-risk sibs.
- If the pathogenic variant(s) in the family are known, molecular genetic testing can be used to clarify the genetic status of at-risk sibs.

See Genetic Counseling for issues related to testing of at-risk relatives for genetic counseling purposes.

Pregnancy Management

Pregnancy is possible in females with FA, whether or not they have undergone hematopoietic stem cell transplantation [Dalle et al 2004, Nabhan et al 2010].

Pregnancy needs to be managed by a high-risk maternal fetal obstetrician along with a hematologist.

Therapies Under Investigation

Previous clinical trials failed to accomplish permanent gene correction of stem cells; current work is focusing on development of novel vector and delivery strategies [Tolar et al 2011]. The first FA lentiviral gene therapy trial led by the University of Washington/Fred Hutchinson Cancer Research Center is now open [Becker et al 2010]. Dr Juan Bueren has an open trial of a hematopoietic stem cell mobilization in Madrid, Spain and plans to have their *FANCA* gene therapy trial opened soon.

A Phase I study of the antioxidant quercetin in children with Fanconi anemia is currently underway at Cincinnati Children's Hospital Medical Center.

Search [ClinicalTrials.gov](https://clinicaltrials.gov) in the US and [EU Clinical Trials Register](https://clinicaltrialsregister.eu) in Europe for access to information on clinical studies for a wide range of diseases and conditions.

Genetic Counseling

Genetic counseling is the process of providing individuals and families with information on the nature, inheritance, and implications of genetic disorders to help them make informed medical and personal decisions. The following section deals with genetic risk assessment and the use of family history and genetic testing to clarify genetic status for family members. This section is not meant to address all personal, cultural, or ethical issues that individuals may face or to substitute for consultation with a genetics professional. —ED.

Mode of Inheritance

Fanconi anemia (FA) is inherited in an autosomal recessive manner, an autosomal dominant manner (*RAD51*-related FA), or an X-linked manner (*FANCB*-related FA).

Autosomal Recessive FA

Risk to Family Members

Parents of a proband

- The parents of a child with autosomal recessive FA are obligate heterozygotes (i.e., carriers of one FA-related pathogenic variant).

- Heterozygotes (carriers) are not at risk for FA. However, heterozygous mutation of a subset of FA-related genes (e.g., *BRCA2*, *PALB2*, and *BRIP1*) is associated with an increased risk for breast and other cancers [Seal et al 2006, Berwick et al 2007, Rahman et al 2007] (see [Hereditary Breast and Ovarian Cancer](#)).

Sibs of a proband

- At conception, each sib of an individual with autosomal recessive FA has a 25% chance of being affected, a 50% chance of being an asymptomatic carrier, and a 25% chance of being unaffected and not a carrier.
- Heterozygotes (carriers) are not at risk for FA. However, heterozygous mutation of a subset of FA-related genes (e.g., *BRCA2*, *PALB2*, and *BRIP1*) is associated with an increased risk for breast and other cancers [Seal et al 2006, Berwick et al 2007, Rahman et al 2007] (see [Hereditary Breast and Ovarian Cancer](#)).

Offspring of a proband. The offspring of an individual with autosomal recessive FA are obligate heterozygotes (carriers) for a pathogenic variant in an FA-related gene.

Other family members of a proband. Each sib of the proband's parents is at a 50% risk of being a carrier for a pathogenic variant in an FA-related gene.

Carrier (Heterozygote) Detection

Carrier testing for at-risk relatives requires prior identification of the FA-related pathogenic variants in the family.

Carriers of FA cannot be detected by the DEB/MMC test.

Autosomal Dominant FA – Risk to Family Members

Parents of a proband

- To date, reported probands with *RAD51*-related FA have the disorder as a result of a *de novo* *RAD51* pathogenic variant.
- Recommendations for the evaluation of parents of a proband with an apparent *de novo* pathogenic variant include molecular genetic testing.

Sibs of a proband. All affected individuals reported to date have had a *de novo* *RAD51*-related FA pathogenic variant, suggesting a low risk to sibs. However, because of the theoretic possibility of germline mosaicism in a parent, the risk is presumed to be greater than in the general population.

Offspring of a proband. Each child of an individual with *RAD51*-related FA is presumed to have a 50% chance of inheriting the pathogenic variant. However, only one individual with *RAD51*-related FA has reached adulthood and no offspring have been reported.

Other family members. Given that all probands with *RAD51*-related FA reported to date have the disorder as a result of a *de novo* *RAD51* pathogenic variant, the risk to other family members is presumed to be low.

X-Linked FA

Risk to Family Members

Parents of a male proband

- The father of a male with X-linked FA will not have the disorder nor will he be hemizygous for the *FANCB* pathogenic variant; therefore, he does not require further evaluation/testing.

- In a family with more than one affected individual, the mother of an affected male is an obligate heterozygote (carrier).

Note: If a woman has more than one affected son and if the pathogenic variant cannot be detected in her leukocyte DNA, she has germline mosaicism.

- If a male is the only affected family member (i.e., a simplex case), the mother may be a carrier or the affected male may have a *de novo* *FANCB* pathogenic variant, in which case the mother is not a carrier.

Sibs of a male proband. The risk to sibs depends on the genetic status of the mother:

- If the mother of the proband has a *FANCB* pathogenic variant, the chance of transmitting it in each pregnancy is 50%. Male sibs who inherit the pathogenic variant will be affected; female sibs who inherit the pathogenic variant will be carriers and will usually not be affected.
- If the proband represents a simplex case (i.e., a single occurrence in a family) and if the *FANCB* pathogenic variant cannot be detected in the leukocyte DNA of the mother, the risk to sibs is low but greater than that of the general population because of the possibility of maternal germline mosaicism.

Offspring of a male proband

- Affected males transmit the *FANCB* pathogenic variant to all of their daughters (who will be carriers and will usually not be affected) and none of their sons.
- To date, no male with FA-B has been old enough to have children; they may also be infertile, as are many males with FA.

Other family members of a proband. The proband's maternal aunts may be at risk of being heterozygotes (carriers) for the *FANCB* pathogenic variant and the aunt's offspring, depending on their gender, may be at risk of being carriers or of being affected.

Note: Molecular genetic testing may be able to identify the family member in whom a *de novo* pathogenic variant arose – information that could help determine genetic risk status of the extended family.

Heterozygote (Carrier) Detection

Carrier testing for at-risk female relatives requires prior identification of the *FANCB* pathogenic variant in the family.

Carriers of FA cannot be detected by the DEB/MMC test.

Related Genetic Counseling Issues

See Management, Evaluation of Relatives at Risk for information on evaluating at-risk relatives for the purpose of early diagnosis and treatment.

Family planning

- The optimal time for determination of genetic risk, clarification of genetic status, and discussion of the availability of prenatal testing is before pregnancy.
- It is appropriate to offer genetic counseling (including discussion of potential risks to offspring and reproductive options) to young adults who are affected, are carriers, or are at risk of being carriers.

DNA banking is the storage of DNA (typically extracted from white blood cells) for possible future use. Because it is likely that testing methodology and our understanding of genes, allelic variants, and diseases will improve in the future, consideration should be given to banking DNA of affected individuals.

Prenatal Testing and Preimplantation Genetic Diagnosis

Molecular genetic testing. Once the FA-related pathogenic variant(s) have been identified in an affected family member, prenatal testing for a pregnancy at increased risk and preimplantation genetic diagnosis for FA are possible. Preimplantation genetic diagnosis has successfully identified at-risk embryos as unaffected with FA and HLA-matched to affected sibs [Kahraman et al 2014].

Chromosome breakage. Prenatal testing is also possible for pregnancies at increased risk for FA by performing cytogenetic testing in the presence of DEB/MMC to evaluate for increased chromosome breakage in fetal cells obtained by chorionic villus sampling (CVS) or amniocentesis; however, if the pathogenic variants are known in the family, molecular genetic testing is the method of choice for prenatal diagnosis.

Fetal ultrasound evaluation. Ultrasound examination can be used to evaluate for fetal anomalies consistent with FA. However, ultrasound examination is not a diagnostic test for FA. Many congenital anomalies characteristic of FA may not be detectable by ultrasound examination, and those that can be seen may be associated with diagnoses other than FA.

Resources

GeneReviews staff has selected the following disease-specific and/or umbrella support organizations and/or registries for the benefit of individuals with this disorder and their families. GeneReviews is not responsible for the information provided by other organizations. For information on selection criteria, click [here](#).

- **Fanconi Anemia Cell Repository**

Department of Medical and Molecular Genetics
3181 Southwest Sam Jackson Park Road L103
Oregon Health & Science University
Portland OR 97201

Phone: 503-494-6888

- **Fanconi Anemia Research Fund, Inc. (FARF)**

1801 Willamette Street
Suite 200
Eugene OR 97401

Phone: 888-326-2664 (Toll-free Family Support Line); 541-687-4658

Fax: 541-687-0548

Email: info@fanconi.org

www.fanconi.org

- **International Fanconi Anemia Registry (IFAR)**

The Rockefeller University
1230 York Avenue
New York NY 10065

Phone: 212-327-8862

Fax: 212-327-8262

Email: auerbac@rockefeller.edu

International Fanconi Anemia Registry (IFAR)

- **National Cancer Institute Inherited Bone Marrow Failure Syndromes (IBMFS) Cohort Registry**

Phone: 800-518-8474

Email: NCI.IBMFS@westat.com

www.marrowsfailure.cancer.gov

- **Prospective Registry of MultiPlex Testing (PROMPT)**

PROMPT is an online research registry for patients and their families that helps researchers answer the question: “How do genetic variants affect your cancer risk?”

PROMPT

Molecular Genetics

Information in the Molecular Genetics and OMIM tables may differ from that elsewhere in the GeneReview: tables may contain more recent information. —ED.

Table A. Fanconi Anemia: Genes and Databases

Gene	Chromosome Locus	Protein	Locus-Specific Databases	HGMD	ClinVar
<i>BRCA2</i>	13q13.1	Breast cancer type 2 susceptibility protein	BRCA2 homepage - LOVD Database of BRCA1 and BRCA2 sequence variants that have been clinically reclassified using a quantitative integrated evaluation Breast Cancer Information Core (BRCA2) Fanconi Anaemia Mutation Database (FANCD1 - BRCA2) BRCA2 @ ZAC-GGM	BRCA2	BRCA2
<i>BRIP1</i>	17q23.2	Fanconi anemia group J protein	BRIP1 @ LOVD Fanconi Anaemia Mutation Database (FANCI - BRIP1)	BRIP1	BRIP1
<i>ERCC4</i>	16p13.12	DNA repair endonuclease XPF	ERCC4 database	ERCC4	ERCC4
<i>FANCA</i>	16q24.3	Fanconi anemia group A protein	Fanconi Anemia Mutation Database (FANCA)	FANCA	FANCA
<i>FANCB</i>	Xp22.2	Fanconi anemia group B protein	FANCB @ LOVD Fanconi Anaemia Mutation Database (FANCB)	FANCB	FANCB
<i>FANCC</i>	9q22.32	Fanconi anemia group C protein	Fanconi Anemia Mutation Database (FANCC)	FANCC	FANCC
<i>FANCD2</i>	3p25.3	Fanconi anemia group D2 protein	Fanconi Anaemia Mutation Database (FANCD2)	FANCD2	FANCD2
<i>FANCE</i>	6p21.31	Fanconi anemia group E protein	Fanconi Anaemia Mutation Database (FANCE)	FANCE	FANCE
<i>FANCF</i>	11p14.3	Fanconi anemia group F protein	Fanconi Anaemia Mutation Database (FANCF)	FANCF	FANCF
<i>FANCG</i>	9p13.3	Fanconi anemia group G protein	Fanconi Anaemia Mutation Database (FANCG)	FANCG	FANCG

Table A. continued from previous page.

<i>FANCI</i>	15q26.1	Fanconi anemia group I protein	Fanconi Anemia Mutation Database (FANCI)	FANCI	FANCI
<i>FANCL</i>	2p16.1	E3 ubiquitin-protein ligase FANCL	Fanconi Anaemia Mutation Database (FANCL)	FANCL	FANCL
<i>FANCM</i>	14q21.2	Fanconi anemia group M protein	Fanconi Anaemia Mutation Database (FANCM)	FANCM	FANCM
<i>MAD2L2</i>	1p36.22	Mitotic spindle assembly checkpoint protein MAD2B		MAD2L2	MAD2L2
<i>PALB2</i>	16p12.2	Partner and localizer of BRCA2	PALB2 database	PALB2	PALB2
<i>RAD51</i>	15q15.1	DNA repair protein RAD51 homolog 1	RAD51 database	RAD51	RAD51
<i>RAD51C</i>	17q22	DNA repair protein RAD51 homolog 3	RAD51C @ LOVD	RAD51C	RAD51C
<i>RFWD3</i>	16q23.1	E3 ubiquitin-protein ligase RFWD3		RFWD3	RFWD3
<i>SLX4</i>	16p13.3	Structure-specific endonuclease subunit SLX4	SLX4 @ LOVD	SLX4	SLX4
<i>UBE2T</i>	1q32.1	Ubiquitin-conjugating enzyme E2 T		UBE2T	UBE2T
<i>XRCC2</i>	7q36.1	DNA repair protein XRCC2	XRCC2 @ LOVD	XRCC2	XRCC2

Data are compiled from the following standard references: gene from [HGNC](#); chromosome locus from [OMIM](#); protein from [UniProt](#). For a description of databases (Locus Specific, HGMD, ClinVar) to which links are provided, click [here](#).

Table B. OMIM Entries for Fanconi Anemia (View All in OMIM)

133520	EXCISION REPAIR, COMPLEMENTING DEFECTIVE, IN CHINESE HAMSTER, 4; ERCC4
179617	RAD51 RECOMBINASE; RAD51
227645	FANCONI ANEMIA, COMPLEMENTATION GROUP C; FANCC
227646	FANCONI ANEMIA, COMPLEMENTATION GROUP D2; FANCD2
227650	FANCONI ANEMIA, COMPLEMENTATION GROUP A; FANCA
300514	FANCONI ANEMIA, COMPLEMENTATION GROUP B; FANCB
300515	FANCB GENE; FANCB
600185	BRCA2 GENE; BRCA2
600375	X-RAY REPAIR, COMPLEMENTING DEFECTIVE, IN CHINESE HAMSTER, 2; XRCC2
600901	FANCONI ANEMIA, COMPLEMENTATION GROUP E; FANCE
602774	RAD51 PARALOG C; RAD51C
602956	FANCG GENE; FANCG
603467	FANCONI ANEMIA, COMPLEMENTATION GROUP F; FANCF
604094	MITOTIC ARREST-DEFICIENT 2 LIKE 2; MAD2L2
605724	FANCONI ANEMIA, COMPLEMENTATION GROUP D1; FANCD1
605882	BRCA1-INTERACTING PROTEIN 1; BRIP1
607139	FANCA GENE; FANCA
608111	FANCL GENE; FANCL
609053	FANCONI ANEMIA, COMPLEMENTATION GROUP I; FANCI

Table B. continued from previous page.

609054	FANCONI ANEMIA, COMPLEMENTATION GROUP J; FANCI
609644	FANCM GENE; FANCM
610355	PARTNER AND LOCALIZER OF BRCA2; PALB2
610538	UBIQUITIN-CONJUGATING ENZYME E2T; UBE2T
610832	FANCONI ANEMIA, COMPLEMENTATION GROUP N; FANCN
611360	FANCI GENE; FANCI
613278	SLX4 STRUCTURE-SPECIFIC ENDONUCLEASE SUBUNIT; SLX4
613390	FANCONI ANEMIA, COMPLEMENTATION GROUP O; FANCO
613897	FANCF GENE; FANCF
613899	FANCC GENE; FANCC
613951	FANCONI ANEMIA, COMPLEMENTATION GROUP P; FANCP
613976	FANCE GENE; FANCE
613984	FANCD2 GENE; FANCD2
614082	FANCONI ANEMIA, COMPLEMENTATION GROUP G; FANCG
614083	FANCONI ANEMIA, COMPLEMENTATION GROUP L; FANCL
614087	FANCONI ANEMIA, COMPLEMENTATION GROUP M; FANCM
614151	RING FINGER AND WD REPEAT DOMAINS-CONTAINING PROTEIN 3; RFWD3
615272	FANCONI ANEMIA, COMPLEMENTATION GROUP Q; FANCQ
616435	FANCONI ANEMIA, COMPLEMENTATION GROUP T; FANCT
617243	FANCONI ANEMIA, COMPLEMENTATION GROUP V; FANCV
617244	FANCONI ANEMIA, COMPLEMENTATION GROUP R; FANCR
617247	FANCONI ANEMIA, COMPLEMENTATION GROUP U; FANCU
617784	FANCONI ANEMIA, COMPLEMENTATION GROUP W; FANCW

See Table A for gene and protein names.

For a table of Fanconi anemia-associated proteins and their function, click [here](#) and download pdf [Wang & Smogorzewska 2015].

Molecular Pathogenesis

At least twenty genes that are involved in Fanconi anemia (FA) and also account for each of the phenotypic complementation groups have been identified. The proteins encoded by the FA-related genes are considered to work together in a common pathway/network called "the FA pathway" or "the FA-BRCA pathway/network," which regulates cellular resistance to DNA cross-linking agents [Taniguchi & D'Andrea 2006, D'Andrea 2010, Deans & West 2011, Kee & D'Andrea 2012]. Disruption of this pathway leads to the common cellular and clinical abnormalities observed in FA [D'Andrea 2010, Nakanishi et al 2011, Williams et al 2011b, Crossan & Patel 2012, Kim & D'Andrea 2012].

Eight of the FA proteins (FANCA, FANCB, FANCC, FANCE, FANCF, FANCG, FANCL, and FANCM), along with proteins FAAP24 [Ciccio et al 2007] and FAAP100 [Ling et al 2007] are assembled in a nuclear complex (FA core complex). This complex is a multisubunit ubiquitin ligase complex; monoubiquitination of two FA proteins (FANCD2 and FANCI) depends on the FA core complex [Garcia-Higuera et al 2001, Smogorzewska et al 2007]. In response to DNA damage or in S phase of the cell cycle, this FA core complex activates the

monoubiquitination of the FANCD2 and FANCI proteins. Monoubiquitinated FANCD2 and monoubiquitinated FANCI are translocated to nuclear foci containing the proteins BRCA1, BRCA2, PALB2, and RAD51. FANCI shares sequence similarity with FANCD2; together they form a protein complex (ID complex) [Smogorzewska et al 2007]. Monoubiquitination of FANCD2 and FANCI is interdependent [Smogorzewska et al 2007]. A nuclease, FAN1, has been shown to bind to monoubiquitinated FANCD2, which directs its enzymatic activity [Huang & D'Andrea 2010]. A cell-free system has been used to recapitulate cross-link repair in vitro [Knipscheer et al 2009].

One of the components of the FA core complex, FANCL, has a PHD (plant homeodomain) finger (a variant RING finger) domain with ubiquitin ligase activity [Meetei et al 2003a]. FANCL associates through its PHD/RING finger domain with UBE2T, a ubiquitin conjugating enzyme (E2), which is also required for FANCD2 monoubiquitination [Machida et al 2006]. Recombinant FANCL, the E2 UBE2T, FANCD2, FANCI, and FANCE recapitulate the monoubiquitination reaction in vitro [Alpi et al 2008].

Another component of the FA core complex, FANCM, is homologous to the archaeal DNA helicase/nuclease known as HEF. FANCM has DNA helicase motifs and a degenerate nuclease motif and exhibits DNA-stimulated ATPase activity and DNA translocase activity [Meetei et al 2005]. A FANCM-interacting protein, FAAP24, preferentially binds to single-stranded DNA and branched DNA structures [Ciccio et al 2007]. Therefore, it has been speculated that FANCM DNA translocase activity could play an important role in displacing the FA core complex along the DNA, allowing DNA damage recognition, or that FAAP24 may play a role in targeting the FA core complex to abnormal, branched DNA structures. This complex is thought to be responsible for a replication-associated checkpoint response involving RPA [Huang et al 2010] and the BLM helicase [Deans & West 2009]. As is true for multiple FA proteins, the complex appears to be regulated through ATR [Collis et al 2008].

Furthermore, the FA core complex forms a larger complex with BLM, RPA, and topoisomerase III α , called BRAFT (BLM, RPA, FA, and topoisomerase III α) [Meetei et al 2003b] in a further link to [Bloom syndrome](#). FANCM is found in both separable complexes: the FA core complex as well as the BLM complex [Deans & West 2009].

A DNA damage-activated signaling kinase, ATR, a single-strand DNA-binding protein complex, RPA, and an ATR-associated protein, HCLK2, are required for DNA damage-inducible monoubiquitination and foci formation of FANCD2 [Andreassen et al 2004, Collis et al 2007]. BRCA1 [Garcia-Higuera et al 2001, Vandenberg et al 2003] and histone H2AX [Bogliolo et al 2007] are required for DNA damage-inducible foci formation of FANCD2, but not for monoubiquitination of FANCD2. These factors are considered to be upstream positive regulators of the FA pathway. ATR has been shown to directly phosphorylate FANCA and indirectly phosphorylate FANCD2 through CHK1 [Zhi et al 2009]. As described, ATR is also necessary for assembly of FANCM and recruitment of RPA at the ICL-induced checkpoint [Collis et al 2008].

BRCA2 (previously *FANCD1*) is a tumor suppressor that confers breast cancer susceptibility [Howlett et al 2002] and has a distinct clinical phenotype [Wagner et al 2004, Alter et al 2007, Myers et al 2012]. *BRCA2* protein stability and localization is regulated by PALB2 (partner and localizer of *BRCA2*) [Xia et al 2006] encoded by *PALB2* (previously *FANCN*), another breast cancer susceptibility gene [Rahman et al 2007]. Another breast cancer susceptibility gene [Seal et al 2006], *BRIP1* (previously *BACH1* or *FANCF*) [Cantor et al 2001], is also associated with FA [Levitus et al 2005, Levrin et al 2005, Litman et al 2005]. *BRCA2*, *PALB2*, and *BRIP1* are not required for FANCD2 protein monoubiquitination or FANCD2 nuclear foci formation, but are still required for cellular resistance to MMC or DEB. *BRCA2* has been found to act in multiple subcomplexes of FA proteins, including FANCG and FANCD2 [Wilson et al 2010], suggesting that the notion of acting downstream of FANCD2 monoubiquitination may be too simplistic. Phosphorylation of FANCD2 by CHK1 has been shown to be necessary for interaction with *BRCA2* [Zhi et al 2009]. *BRIP1* and FANCD2 have also been shown to be functionally linked in foci formation [Zhang et al 2010].

USP1 is a deubiquitinating enzyme that removes ubiquitin from monoubiquitinated FANCD2, and negatively regulates the FA pathway along with its coactivator UAF1 [Nijman et al 2005, Cohn et al 2007]. USP1 also removes ubiquitin from monoubiquitinated PCNA (*proliferating cell nuclear antigen*) [Huang et al 2006]. This may not be coincidental, since FANCD2 and PCNA have been shown to bind [Howlett et al 2009]. The deubiquitination event has been shown to be vital for FA function [Oestergaard et al 2007]. Hematopoietic defects have been noted in knockout mice [Parmar et al 2010].

In nuclear foci, *FANCD2* colocalizes with *FANCI*, *BRCA1*, *BRCA2*, *PALB2*, *RAD51*, *BLM*, *RPA*, *ATR*, *FANCC*, and *FANCE* [Garcia-Higuera et al 2001, Pace et al 2002, Taniguchi et al 2002a, Andreassen et al 2004, Wang et al 2004, Matsushita et al 2005, Xia et al 2006, Smogorzewska et al 2007]. *FANCD2* also colocalizes partially with *BRIP1* [Litman et al 2005] and *NBS1* [Nakanishi et al 2002]. All of these factors are required for cellular resistance to DNA cross-linking agents and are considered to work together to repair interstrand DNA cross-links, although the precise mechanism is not understood. Recently, pathogenic variants in *RAD51C* have been detected in several FA-like cases in a consanguineous family, also associated with breast and ovarian cancer susceptibility [Meindl et al 2010, Vaz et al 2010].

Among FA proteins, *BRCA2* has a clear role in regulating homologous recombination by controlling the activity of *RAD51*, the eukaryotic homolog of bacterial *RecA* [Davies et al 2001, Moynahan et al 2001]. *PALB2* regulates *BRCA2* stability and localization in nuclear structures (chromatin and nuclear matrix) and, thus, is required for homologous recombination [Xia et al 2006]. The FA core complex, *FANCD2*, *FANCI* [Smogorzewska et al 2007], and *BRIP1* [Litman et al 2005] are also reported to be required for efficient homologous recombination, although conflicting reports exist [Taniguchi & D'Andrea 2006].

FANCD2 protein is also phosphorylated by the ataxia-telangiectasia kinase, *ATM*, in a process that regulates a radiation-induced S phase checkpoint [Taniguchi et al 2002b, Ho et al 2006]. While required for resistance to ionizing radiation, this phosphorylation event is dispensable for cross-linker resistance, implying a separation of or dual function for *FANCD2*. *FANCD2* appears to be phosphorylated by *CHK1*, which is downstream of *ATR*, at serine 331 in a manner that results in activation by cross-links [Zhi et al 2009].

Importantly, a number of studies have shown defects in the FA-BRCA pathway to be implicated in cancer:

- Individuals with FA are susceptible to both leukemia and solid tumors [Alter 2003].
- *Fancd2*, *Fanca*, or *Fancc* knockout mice develop tumors [Houghtaling et al 2003, Wong et al 2003, Carreau 2004].
- Inactivation of the FA pathway by somatic methylation of *FANCF* has been found in a wide variety of human cancers (ovarian, breast, non-small cell lung, cervical, testicular, and head and neck squamous cell cancers) in the general population [Olopade & Wei 2003, Taniguchi et al 2003, Marsit et al 2004, Narayan et al 2004, Wang et al 2006].
- Inherited and somatic variants of *FANCC* and *FANCG* are present in a subset of young-onset pancreatic cancers [van der Heijden et al 2003].
- *BRCA1* and *BRCA2* are well-known tumor suppressor genes responsible for familial breast/ovarian cancer [Turner et al 2004].
- Pathogenic truncating variants in the FA-related genes *BRIP1* and *PALB2* are breast cancer susceptibility alleles [Seal et al 2006, Erkkö et al 2007, Rahman et al 2007, Tischkowitz et al 2007].

Amelioration of FA pathology has been implicated in reports of downregulation of elements of the non-homologous end-joining pathway [Adamo et al 2010]. These data propose that much of FA pathophysiology results from the unfettered work of NHEJ promoting inaccurate repair. On the other hand, FA involvement in homologous recombinatorial repair has been well established in interactions with *BRCA1*, *BRCA2*, and *RAD51C*. *FANCD2* has also been shown to interact with *PCNA* and *pol K*, suggesting that translesion synthesis,

a variant of homologous recombination, may be the most direct function of FA proteins in bypass of the lesion [Ho & Schärer 2010, Song et al 2010].

For reviews of the molecular biology of FA, see Nakanishi et al [2011], Williams et al [2011b], Crossan & Patel [2012], Kim & D'Andrea [2012], and Huang et al [2014].

BRCA2 (previously FANCD1)

Gene structure. *BRCA2* has 27 exons (NM_000059.3).

Pathogenic variants. See Table A.

Normal gene product. The breast cancer type 2 susceptibility protein (BRCA2) has 3418 amino acids. BRCA2 regulates homologous recombination repair through control of RAD51 recombinase [Davies et al 2001, Moynahan et al 2001]. Other functions of BRCA2 include stabilizing stalled replication forks and regulating cytokinesis [Daniels et al 2004]. BRCA2 works in concert with RAD51 to orchestrate the repair of double-strand break ends, prompting the nucleation of each end by RAD51. BRCA2 has been found in various subcomplexes with other FA proteins, including FANCG, FANCD2, and PALB2 [Xia et al 2007, Wilson et al 2010].

Abnormal gene product. See Molecular Pathogenesis.

BRIP1 (previously FANCI or BACH1)

Gene structure. *BRIP1* (NM_032043.2) has 20 exons.

Pathogenic variants. See Table A.

Normal gene product. The Fanconi anemia group J protein (BRIP1 or FANCI) has 1249 amino acids and is a DNA-dependent ATPase and a 5'-to-3' DNA helicase (DEAH helicase) that binds directly to the BRCT domain of BRCA1 [Cantor et al 2001, Chen et al 2014, Zou et al 2014]. FANCI acetylation regulates DNA damage response [Xie et al 2012], and it contains the seven helicase-specific motifs and C-terminal extension, which has 39% homology with synaptonemal complex protein 1, a major component of the transverse filaments of developing meiotic chromosomes [Cantor et al 2001, Wu et al 2012, Brosh & Cantor 2014, Sommers et al 2014]. FANCI engages stable G4/G-quadruplex structures [Sarkies et al 2012, Castillo Bosch et al 2014], promotes stability of FANCD2/FANCI [Zhang et al 2010, Clark et al 2015], and its helicase domain clearly is important for FA pathway function [Wu & Brosh 2009, Guo et al 2014]. FANCI also appears to interdigitate with the mismatch repair pathway in binding to MLH1 [Cantor & Xie 2010] and centrosome amplification [Zou et al 2013].

Abnormal gene product. See Molecular Pathogenesis.

FANCA

Gene structure. The longest transcript variant NM_000135.2 has 43 exons.

Pathogenic variants. The pathogenic variants of *FANCA* are numerous and highly variable among families [Levrán et al 1997, Morgan et al 1999, Wijker et al 1999]. A small percentage of families share the pathogenic variants p.Phe1263del and p.Val372AlafsTer42; the latter is found in affected individuals of northern European ancestry. See Table A.

Table 3. Selected *FANCA* Pathogenic Variants

DNA Nucleotide Change	Predicted Protein Change	Reference Sequences
c.1115_1118del	p.Val372AlafsTer42	NM_000135.2
c.3788_3790del	p.Phe1263del	NP_000126.2

Variants listed in the table have been provided by the authors. *GeneReviews* staff have not independently verified the classification of variants.

GeneReviews follows the standard naming conventions of the Human Genome Variation Society (varnomen.hgvs.org). See [Quick Reference](#) for an explanation of nomenclature.

Normal gene product. *FANCA* encoded by [NM_000135.2](#) has 1455 amino acids and is a component of the FA core complex. *FANCA* contains two overlapping bipartite nuclear localization signals (NLS), five functional leucine-rich nuclear export sequences, and a partial leucine zipper sequence [Fanconi Anaemia/Breast Cancer Consortium 1996, Lo Ten Foe et al 1996, Ferrer et al 2005]. The nuclear export of *FANCA* is regulated in a CRM1-dependent manner [Ferrer et al 2005]. *FANCA* is a phosphoprotein, a client of Hsp90 [Oda et al 2007]; it is phosphorylated by the ATR kinase at serine 1449 [Collins et al 2009]. Reports show that *FANCA* interacts with FA binding protein FAAP20 [Ali et al 2012, Leung et al 2012], is co-regulated with BRCA1 [Haitjema et al 2014], and its deletion dysregulates murine T regulatory cells [Du et al 2014]. With *FANCC* it modulates the toll-like receptor pathway [Garbati et al 2013] and – critically – safeguards cell cycle integrity in hematopoietic cells [Abdul-Sater et al 2015].

Abnormal gene product. See Molecular Pathogenesis.

FANCB

Gene structure. *FANCB* has ten exons with the translation start in exon 3 ([NM_001018113.1](#)).

Pathogenic variants. See Table A.

Normal gene product. *FANCB* comprises 853 amino acids; some sequences have 859 residues, depending on the initiating methionine. *FANCB* is a component of the FA core complex and contains a putative bipartite NLS [Meetei et al 2004]. *FANCB* is relevant to normal development – as is evidenced by its dysfunction being linked to VACTERL association [McCauley et al 2011, Umaña et al 2011].

Abnormal gene product. See Molecular Pathogenesis.

FANCC

Gene structure. *FANCC* has 15 exons.

Pathogenic variants. Three common pathogenic variants in *FANCC* have been identified (c.456+4A>T, c.1642C>T, and c.67delG) [Whitney et al 1993]. See Table A.

Table 4. Selected *FANCC* Pathogenic Variants

DNA Nucleotide Change (Alias ¹)	Predicted Protein Change	Reference Sequences
c.456+4A>T ^{2, 3} (IVS4+4A>T)	--	NM_000136.2 NP_000127.2
c.37C>T ^{3, 4}	p.Gln13Ter	
c.67delG ^{4, 5} (322delG)	p.Asp23IlefsTer23	
c.1642C>T ^{4, 5}	p.Arg548Ter	
c.1661T>C ³	p.Leu554Pro	

Variants listed in the table have been provided by the authors. *GeneReviews* staff have not independently verified the classification of variants.

GeneReviews follows the standard naming conventions of the Human Genome Variation Society (varnomen.hgvs.org). See [Quick Reference](#) for an explanation of nomenclature.

1. Variant designation that does not conform to current naming conventions
2. Found primarily in the Ashkenazi Jewish population; but also reported in a Japanese cohort.
3. See Genotype-Phenotype Correlations.
4. Prevalent in individuals of northern European ancestry
5. Found in individuals from southern Italy

Normal gene product. *FANCC* has 558 amino acids. It is a component of the FA core complex, but localizes to both the nucleus and the cytoplasm [Yamashita et al 1994]. Some functions of *FANCC* outside of the FA core complex have also been proposed [Fagerlie et al 2004]. Reports suggest that STAT1 binds to *FANCC* to modulate JAK-STAT signaling and to protect cells from interferon gamma toxic effects [Pang et al 2000, Fagerlie et al 2004]. Further, *FANCC* binds to microtubule-associated protein stathmin-1 to regulate cytokinesis [Magron et al 2015].

Abnormal gene product. See Molecular Pathogenesis.

FANCD2

Gene structure. *FANCD2* has two transcript variants. Variant 1 (NM_033084.3) has 43 exons; variant 2 (NM_001018115.1) has 44 exons and an alternate 3' coding sequence resulting in a shorter and distinct C-terminus. *FANCD2* protein encoded by variant 2 (exon 44 form) is the functional *FANCD2* protein, while the other transcript variant encodes a non-functional protein [Montes de Oca et al 2005].

Pathogenic variants. See Table A.

Normal gene product. The functional *FANCD2* (NP_001018125.1) has 1451 amino acids (isoform b) and shares sequence similarity with *FANCI*. *FANCD2* and *FANCI* form a protein complex (ID complex). *FANCD2* can be monoubiquitinated on lysine 561 in an FA core complex-, UBE2T- (also known as *FANCT* [Hira et al 2015, Rickman et al 2015, Virts et al 2015]), and *FANCI*-dependent manner [Sato et al 2012, Rajendra et al 2014]. Reports illuminated the structure of *FANCI-FANCD2* complex [Joo et al 2011], its coordinate nuclear targeting [Boisvert et al 2013], and its RAD18-mediated chromatin loading [Song et al 2010, Williams et al 2011a]. Monoubiquitinated *FANCD2* is translocated to chromatin fraction – recruited by UHRF1 [Liang et al 2015] – to form nuclear foci with *FANCI*, BRCA1 (*FANCS* [Sawyer et al 2015]), BRCA2, RAD51, and other partners. *FANCD2* can be phosphorylated by ATM [Taniguchi et al 2002b, Ho et al 2006] and by ATR [Andreassen et al 2004, Pichierri & Rosselli 2004] in response to DNA damage. *FANCD2* is phosphorylated at serine 331 by CHK1 in a manner that is required for binding to BRCA2 [Zhi et al 2009]. Important studies have shown *FANCD2* to have a function in resection of ends surrounding a cross-link in the repair process [Knipscheer et al 2009], in replication fork processes [Zhu et al 2015b], binding to mini-chromosome maintenance proteins in control of

replisome [Lossaint et al 2013]. A nuclease, FAN1, has been demonstrated to bind to FANCD2 [Huang & D'Andrea 2010, Kratz et al 2010, MacKay et al 2010], and to promote DNA interstrand cross-link repair [Smogorzewska et al 2010, Yoshikiyo et al 2010].

Abnormal gene product. See Molecular Pathogenesis.

FANCE

Gene structure. *FANCE* has 14 exons ([NM_021922.2](#)).

Pathogenic variants. See Table A.

Normal gene product. FANCE has 536 amino acids and is a component of the FA core complex. FANCE directly binds to FANCD2 and recruits it to the FA E3 ligase complex [Polito et al 2014]. FANCE contains two nuclear localization signals (NLS) and its nuclear accumulation depends on FANCC [Léveillé et al 2006]. FANCE has five tandem repeats of a short helical motif (FANC repeats) [Nookala et al 2007]. It functions as a shuttle protein between the FA core complex and FANCD2 in a fashion dependent on phosphorylation [Wang et al 2007], with functionality dependent on alternative splicing [Bouffard et al 2015].

Abnormal gene product. See Molecular Pathogenesis.

FANCF

Gene structure. *FANCF* has a single exon ([NM_022725.2](#)).

Pathogenic variants. See Table A.

Normal gene product. FANCF has 374 amino acids and is a component of the FA core complex. FANCF acts as a flexible adaptor protein required for the assembly of the FA core complex [Léveillé et al 2004]. Crystallographic studies of the C-terminal domain revealed a helical repeat structure similar to the Cand1 regulator of the Cul1-Rbx1-Skp1-Fbox(Skp2) ubiquitin ligase complex [Kowal et al 2007], and – recently – next-generation sequencing refined genotype-phenotype correlations [Nicchia et al 2015].

Abnormal gene product. See Molecular Pathogenesis.

FANCG

Gene structure. *FANCG* has 14 exons (reference sequence [NM_004629.1](#)).

Pathogenic variants. Pathogenic variants in *FANCG* are highly variable, but more common variant alleles have been described in specific populations: c.307+1G>C (Korean/Japanese), c.925-2A>G (Brazilian), c.1480+1G>C (French Canadian), and p.Gly395TrpfsTer5 [Demuth et al 2000] and p.Trp599ProfsTer49 [Auerbach et al 2003] (northern European). See Table A.

Table 5. Selected *FANCG* Pathogenic Variants

DNA Nucleotide Change (Alias ¹)	Predicted Protein Change	Reference Sequences
c.307+1G>C (IVS3+1G>C)	--	NM_004629.1 NP_004620.1
c.925-2A>G (IVS8-2A>G)	--	
c.1183-1192del (1184-1194del)	p.Glu395TrpfsTer5	
c.1480+1G>C (IVS11+1G>C)	p.Trp599ProfsTer49	
c.1794_1803del		

Variants listed in the table have been provided by the authors. *GeneReviews* staff have not independently verified the classification of variants.

GeneReviews follows the standard naming conventions of the Human Genome Variation Society (varnomen.hgvs.org). See [Quick Reference](#) for an explanation of nomenclature.

1. Variant designation that does not conform to current naming conventions

Normal gene product. *FANCG* has 622 amino acids. It is a component of the FA core complex. *FANCG* has seven tetratricopeptide repeat motifs [Blom et al 2004, Wilson et al 2010]. *FANCG* is a phosphoprotein; serines 383 and 387 on *FANCG* are phosphorylated in M phase, presumably by cdc2 [Mi et al 2004]. These two sites are important for exclusion of *FANCG* from chromatin in mitosis. Phosphorylation of serine 7 of *FANCG* is upregulated after MMC treatment [Qiao et al 2004]. *FANCA* and *FANCG* stabilize each other, and *FANCG*, *BRCA2*, *FANCD2*, and *XRCC3* participate in the same protein complex. This implies multifunctionality of *FANCG* by its presence in the core complex as well as in homologous recombinatorial repair [Wang et al 2007]. *FANCG* further associates with Rap80-*BRCA1* complex in DNA repair mediated by homologous recombination [Zhu et al 2015a], and it is operational in homing of hematopoietic stem cells [Barroca et al 2012].

Abnormal gene product. See Molecular Pathogenesis.

FANCI

Gene structure. *FANCI* has 37 exons (NM_018193.2).

Pathogenic variants. See Table A.

Normal gene product. *FANCI* has 1268 amino acids and shares sequence similarity with *FANCD2*. *FANCD2* and *FANCI* form a protein complex (ID complex). *FANCI* can be monoubiquitinated on lysine 523 in an FA core complex-, UBE2T-, and *FANCD2*-dependent manner. Monoubiquitinated *FANCI* is translocated to nuclear foci and colocalizes with *BRCA1*, *BRCA2*, *RAD51*, *FANCD2* [Boisvert & Howlett 2014, Rajendra et al 2014]. *FANCI* is a phosphoprotein. DNA damage-induced phosphorylation of p.Ser730, p.Thr952, and p.Ser1121 of human *FANCI* can be detected [Smogorzewska et al 2007]. *FANCI* functions in an analogous manner as *FANCD2* and can be analyzed in the same fashion in assays including DNA repair foci, cell survival, and monoubiquitination [Ishiai et al 2008]. Reports illuminated the structure of *FANCI*-*FANCD2* complex [Joo et al 2011], its coordinate nuclear targeting [Boisvert et al 2013], and – remarkably – its *FANCD2*-independent function upstream of FA core complex recruitment [Castella et al 2015].

Abnormal gene product. See Molecular Pathogenesis.

Click [here](#) (pdf) for information on the genes less commonly involved in Fanconi Anemia (from Table 1b).

References

Literature Cited

- Abdul-Sater Z, Cerabona D, Potchanant ES, Sun Z, Enzor R, He Y, Robertson K, Goebel WS, Nalepa G. FANCA safeguards interphase and mitosis during hematopoiesis in vivo. *Exp Hematol.* 2015;43:1031–46.e12. PubMed PMID: 26366677.
- Adamo A, Collis SJ, Adelman CA, Silva N, Horejsi Z, Ward JD, Martinez-Perez E, Boulton SJ, La Volpe A. Preventing nonhomologous end joining suppresses DNA repair defects of Fanconi anemia. *Mol Cell.* 2010;39:25–35. PubMed PMID: 20598602.
- Ali AM, Pradhan A, Singh TR, Du C, Li J, Wahengbam K, Grassman E, Auerbach AD, Pang Q, Meetei AR. FAAP20: a novel ubiquitin-binding FA nuclear core-complex protein required for functional integrity of the FA-BRCA DNA repair pathway. *Blood.* 2012;119:3285–94. PubMed PMID: 22343915.
- Alpi AF, Pace PE, Babu MM, Patel KJ. Mechanistic insight into site-restricted monoubiquitination of FANCD2 by Ube2t, FANCL, and FANCI. *Mol Cell.* 2008;32:767–77. PubMed PMID: 19111657.
- Alter BP. Inherited bone marrow failure syndromes. In: Nathan DG, Orkin SH, Look AT, Ginsburg D, eds. *Nathan and Oski's Hematology of Infancy and Childhood.* 6 ed. Philadelphia, PA: WB Saunders; 2003:280-365.
- Alter BP. The association between FANCD1/BRCA2 mutations and leukaemia. *Br J Haematol.* 2006;133:446–8. PubMed PMID: 16643458.
- Alter BP, Giri N, Savage SA, Peters JA, Loud JT, Leathwood L, Carr AG, Greene MH, Rosenberg PS. Malignancies and survival patterns in the National Cancer Institute inherited bone marrow failure syndromes cohort study. *Br J Haematol.* 2010;150:179–88. PubMed PMID: 20507306.
- Alter BP, Rosenberg PS, Brody LC. Clinical and molecular features associated with biallelic mutations in FANCD1/BRCA2. *J Med Genet.* 2007;44:1–9. PubMed PMID: 16825431.
- Ameziane N, May P, Haitjema A, van de Vrugt HJ, van Rossum-Fikkert SE, Ristic D, Williams GJ, Balk J, Rockx D, Li H, Rooimans MA, Oostra AB, Velleuer E, Dietrich R, Bleijerveld OB, Maarten Altelaar AF, Meijers-Heijboer H, Joenje H, Glusman G, Roach J, Hood L, Galas D, Wyman C, Balling R, den Dunnen J, de Winter JP, Kanaar R, Gelinas R, Dorsman JC. A novel Fanconi anaemia subtype associated with a dominant-negative mutation in RAD51. *Nat Commun.* 2015;6:8829. PubMed PMID: 26681308.
- Andreassen PR, D'Andrea AD, Taniguchi T. ATR couples FANCD2 monoubiquitination to the DNA-damage response. *Genes Dev.* 2004;18:1958–63. PubMed PMID: 15314022.
- Auerbach AD, Greenbaum J, Pujara K, Batish SD, Bitencourt MA, Kokemohr I, Schneider H, Lobitz S, Pasquini R, Giampietro PF, Hanenberg H, Levran O; International Fanconi Anemia Registry. Spectrum of sequence variation in the FANCG gene: an International Fanconi Anemia Registry (IFAR) study. *Hum Mutat.* 2003;21:158–68. PubMed PMID: 12552564.
- Barroca V, Mouthon MA, Lewandowski D, Brunet de la Grange P, Gauthier LR, Pflumio F, Boussin FD, Arwert F, Riou L, Allemand I, Romeo PH, Fouchet P. Impaired functionality and homing of Fancg-deficient hematopoietic stem cells. *Hum Mol Genet.* 2012;21:121–35. PubMed PMID: 21968513.
- Becker PS, Taylor JA, Trobridge GD, Zhao X, Beard BC, Chien S, Adair J, Kohn DB, Wagner JE, Shimamura A, Kiem HP. Preclinical correction of human Fanconi anemia complementation group A bone marrow cells using a safety-modified lentiviral vector. *Gene Ther.* 2010;17:1244–52. PubMed PMID: 20485382.
- Beier R, Maecker-Kolhoff B, Sykora K-W, Chao M, Kratz C, Sauer MG. Minimal antileukaemic treatment followed by reduced-intensity conditioning in three consecutive children with Fanconi anaemia and AML. *Bone Marrow Transplant.* 2015;50:463–4. PubMed PMID: 25531282.
- Berwick M, Satagopan JM, Ben-Porat L, Carlson A, Mah K, Henry R, Diotti R, Milton K, Pujara K, Landers T, Dev Batish S, Morales J, Schindler D, Hanenberg H, Hromas R, Levran O, Auerbach AD. Genetic

- heterogeneity among Fanconi anemia heterozygotes and risk of cancer. *Cancer Res.* 2007;67:9591–6. PubMed PMID: 17909071.
- Blom E, van de Vrugt HJ, de Vries Y, de Winter JP, Arwert F, Joenje H. Multiple TPR motifs characterize the Fanconi anemia FANCG protein. *DNA Repair (Amst).* 2004;3:77–84. PubMed PMID: 14697762.
- Bluteau D, Masliah-Planchon J, Clairmont C, Rousseau A, Ceccaldi R, Dubois d'Enghien C, Bluteau O, Cuccuini W, Gachet S, Peffault de Latour R, Leblanc T, Socié G, Baruchel A, Stoppa-Lyonnet D, D'Andrea AD, Soulier J. Biallelic inactivation of REV7 is associated with Fanconi anemia. *J Clin Invest.* 2016;126:3580–4. PubMed PMID: 27500492.
- Bogliolo M, Lyakhovich A, Callén E, Castellà M, Cappelli E, Ramírez MJ, Creus A, Marcos R, Kalb R, Neveling K, Schindler D, Surrallés J. Histone H2AX and Fanconi anemia FANCD2 function in the same pathway to maintain chromosome stability. *EMBO J.* 2007;26:1340–51. PubMed PMID: 17304220.
- Bogliolo M, Schuster B, Stoepker C, Derkunt B, Su Y, Raams A, Trujillo JP, Minguillón J, Ramírez MJ, Pujol R, Casado JA, Baños R, Rio P, Knies K, Zúñiga S, Benítez J, Bueren JA, Jaspers NG, Schärer OD, de Winter JP, Schindler D, Surrallés J. Mutations in ERCC4, encoding the DNA-repair endonuclease XPF, cause Fanconi anemia. *Am J Hum Genet.* 2013;92:800–6. PubMed PMID: 23623386.
- Bonfim C, Riberio L, Nichele S, Loth G, Kuwahara C, Koliski A, Bitencourt M, Scherer F, Rodrigues L, Pilonetto D, Pasquini R. Excellent outcomes for Fanconi anemia patients undergoing hematopoietic stem cell transplantation (HSCT) without radiation: a single center experience on 103 patients. *Biol Blood Marrow Transplant.* 2015;21:S94a.
- Boisvert RA, Howlett NG. The Fanconi anemia ID2 complex: dueling saxes at the crossroads. *Cell Cycle.* 2014;13:2999–3015. PubMed PMID: 25486561.
- Boisvert RA, Rego MA, Azzinaro PA, Mauro M, Howlett NG. Coordinate nuclear targeting of the FANCD2 and FANCI proteins via a FANCD2 nuclear localization signal. *PLoS One.* 2013;8:e81387. PubMed PMID: 24278431.
- Bouffard F, Plourde K, Bélanger S, Ouellette G, Labrie Y, Durocher F. Analysis of a FANCE Splice Isoform in Regard to DNA Repair. *J Mol Biol.* 2015;427:3056–73. PubMed PMID: 26277624.
- Brosh RM Jr, Cantor SB. Molecular and cellular functions of the FANCI DNA helicase defective in cancer and in Fanconi anemia. *Front Genet.* 2014;5:372. PubMed PMID: 25374583.
- Byrd PJ, Stewart GS, Smith A, Eaton C, Taylor AJ, Guy C, Eringyte I, Fooks P, Last JI, Horsley R, Oliver AW, Janic D, Dokmanovic L, Stankovic T, Taylor AM. A hypomorphic PALB2 allele gives rise to an unusual form of FA-N associated with lymphoid tumour development. *PLoS Genet.* 2016;12:e1005945. PubMed PMID: 26990772.
- Cantor SB, Bell DW, Ganesan S, Kass EM, Drapkin R, Grossman S, Wahrer DC, Sgroi DC, Lane WS, Haber DA, Livingston DM. BACH1, a novel helicase-like protein, interacts directly with BRCA1 and contributes to its DNA repair function. *Cell.* 2001;105:149–60. PubMed PMID: 11301010.
- Cantor SB, Xie J. Assessing the link between BACH1/FANCI and MLH1 in DNA crosslink repair. *Environ Mol Mutagen.* 2010;51:500–7. PubMed PMID: 20658644.
- Carreau M. Not-so-novel phenotypes in the Fanconi anemia group D2 mouse model. *Blood.* 2004;103:2430. PubMed PMID: 14998919.
- Castella M, Jacquemont C, Thompson EL, Yeo JE, Cheung RS, Huang JW, Sobeck A, Hendrickson EA, Taniguchi T. FANCI regulates recruitment of the FA core complex at sites of DNA damage independently of FANCD2. *PLoS Genet.* 2015;11:e1005563. PubMed PMID: 26430909.
- Castillo Bosch P, Segura-Bayona S, Koole W, van Heteren JT, Dewar JM, Tijsterman M, Knipscheer P. FANCI promotes DNA synthesis through G-quadruplex structures. *EMBO J.* 2014;33:2521–33. PubMed PMID: 25193968.

- Chao MM, Kuehl JS, Strauss G, Hanenberg H, Schindler D, Neitzel H, Niemeyer C, Baumann I, von Bernuth H, Rascon J, Nagy M, Zimmermann M, Kratz CP, Ebell W. Outcomes of mismatched and unrelated donor hematopoietic stem cell transplantation in Fanconi anemia conditioned with chemotherapy only. *Ann Hematol.* 2015;94:1311–8. PubMed PMID: 25862235.
- Chandrasekharappa SC, Lach FP, Kimble DC, Kamat A, Teer JK, Donovan FX, Flynn E, Sen SK, Thongthip S, Sanborn E, Smogorzewska A, Auerbach AD, Ostrander EA; NISC Comparative Sequencing Program. Massively parallel sequencing, aCGH, and RNA-Seq technologies provide a comprehensive molecular diagnosis of Fanconi anemia. *Blood.* 2013;121:e138–48. PubMed PMID: 23613520.
- Chaudhury S, Auerbach AD, Kernan NA, Small TN, Prockop SE, Scaradavou A, Heller G, Wolden S, O'Reilly RJ, Boulad F. Fludarabine-based cytoreductive regimen and T-cell-depleted grafts from alternative donors for the treatment of high-risk patients with Fanconi anaemia. *Br J Haematol.* 2008;140:644–55. PubMed PMID: 18302713.
- Chen X, Wilson JB, McChesney P, Williams SA, Kwon Y, Longerich S, Marriott AS, Sung P, Jones NJ, Kupfer GM. The Fanconi anemia proteins FANCD2 and FANCI interact and regulate each other's chromatin localization. *J Biol Chem.* 2014;289:25774–82. PubMed PMID: 25070891.
- Chrzanowska KH, Gregorek H, Dembowska-Bagińska B, Kalina MA, Digweed M. Nijmegen breakage syndrome (NBS). *Orphanet J Rare Dis.* 2012;7:13. PubMed PMID: 22373003.
- Ciccio A, Ling C, Coulthard R, Yan Z, Xue Y, Meetei AR, Laghmani el H, Joenje H, McDonald N, de Winter JP, Wang W, West SC. Identification of FAAP24, a Fanconi anemia core complex protein that interacts with FANCM. *Mol Cell.* 2007;25:331–43. PubMed PMID: 17289582.
- Clark DW, Tripathi K, Dorsman JC, Palle K. FANCI protein is important for the stability of FANCD2/FANCI proteins and protects them from proteasome and caspase-3 dependent degradation. *Oncotarget.* 2015;6:28816–32. PubMed PMID: 26336824.
- Cohn MA, Kowal P, Yang K, Haas W, Huang TT, Gygi SP, D'Andrea AD. A UAF1-containing multisubunit protein complex regulates the Fanconi anemia pathway. *Mol Cell.* 2007;28:786–97. PubMed PMID: 18082604.
- Collins NB, Wilson JB, Bush T, Thomashevski A, Roberts KJ, Jones NJ, Kupfer GM. ATR-dependent phosphorylation of FANCA on serine 1449 after DNA damage is important for FA pathway function. *Blood.* 2009;113:2181–90. PubMed PMID: 19109555.
- Collis SJ, Ciccio A, Deans AJ, Horejsí Z, Martin JS, Maslen SL, Skehel JM, Elledge SJ, West SC, Boulton SJ. FANCM and FAAP24 function in ATR-mediated checkpoint signaling independently of the Fanconi anemia core complex. *Mol Cell.* 2008;32:313–24. PubMed PMID: 18995830.
- Collis SJ, Barber LJ, Clark AJ, Martin JS, Ward JD, Boulton SJ. HCLK2 is essential for the mammalian S-phase checkpoint and impacts on Chk1 stability. *Nat Cell Biol.* 2007;9:391–401. PubMed PMID: 17384638.
- Crossan GP, Patel KJ. The Fanconi anaemia pathway orchestrates incisions at sites of crosslinked DNA. *J Pathol.* 2012;226:326–37. PubMed PMID: 21956823.
- D'Andrea AD. Susceptibility pathways in Fanconi's anemia and breast cancer. *N Engl J Med.* 2010;362:1909–19. PubMed PMID: 20484397.
- Dalle JH, Huot C, Duval M, Rousseau P, Francoeur D, Champagne J, Vachon MF, Champagne MA. Successful pregnancies after bone marrow transplantation for Fanconi anemia. *Bone Marrow Transplant.* 2004;34:1099–100. PubMed PMID: 15489874.
- Daniels MJ, Wang Y, Lee M, Venkitaraman AR. Abnormal cytokinesis in cells deficient in the breast cancer susceptibility protein BRCA2. *Science.* 2004;306:876–9. PubMed PMID: 15375219.

- Davies AA, Masson JY, McIlwraith MJ, Stasiak AZ, Stasiak A, Venkitaraman AR, West SC. Role of BRCA2 in control of the RAD51 recombination and DNA repair protein. *Mol Cell*. 2001;7:273–82. PubMed PMID: 11239456.
- Deans AJ, West SC. FANCM connects the genome instability disorders Bloom's Syndrome and Fanconi Anemia. *Mol Cell*. 2009;36:943–53. PubMed PMID: 20064461.
- Deans AJ, West SC. DNA interstrand crosslink repair and cancer. *Nat Rev Cancer*. 2011;11:467–80. PubMed PMID: 21701511.
- Demuth I, Wlodarski M, Tipping AJ, Morgan NV, de Winter JP, Thiel M, Gräsl S, Schindler D, D'Andrea AD, Altay C, Kayserili H, Zatterale A, Kunze J, Ebell W, Mathew CG, Joenje H, Sperling K, Digweed M. Spectrum of mutations in the Fanconi anaemia group G gene, FANCG/XRCC9. *Eur J Hum Genet*. 2000;8:861–8. PubMed PMID: 11093276.
- Du W, Erden O, Wilson A, Sippl JM, Schick J, Mehta P, Myers KC, Steinbrecher KA, Davies SM, Pang Q. Deletion of *Fancc* or *Fancc2* dysregulates Treg in mice. *Blood*. 2014;123:1938–47. PubMed PMID: 24501220.
- Erkko H, Xia B, Nikkilä J, Schleutker J, Syrjäkoski K, Mannermaa A, Kallioniemi A, Pylkäs K, Karppinen SM, Rapakko K, Miron A, Sheng Q, Li G, Mattila H, Bell DW, Haber DA, Grip M, Reiman M, Jukkola-Vuorinen A, Mustonen A, Kere J, Aaltonen LA, Kosma VM, Kataja V, Soini Y, Drapkin RI, Livingston DM, Winqvist R. A recurrent mutation in PALB2 in Finnish cancer families. *Nature*. 2007;446:316–9. PubMed PMID: 17287723.
- Fagerlie SR, Koretsky T, Torok-Storb B, Bagby GC. Impaired type I IFN-induced Jak/STAT signaling in FA-C cells and abnormal CD4+ Th cell subsets in *Fancc*^{-/-} mice. *J Immunol*. 2004;173:3863–70. PubMed PMID: 15356134.
- Faivre L, Guardiola P, Lewis C, Dokal I, Ebell W, Zatterale A, Altay C, Poole J, Stones D, Kwee ML, van Weel-Sipman M, Havenga C, Morgan N, de Winter J, Digweed M, Savoia A, Pronk J, de Ravel T, Jansen S, Joenje H, Gluckman E, Mathew CG. Association of complementation group and mutation type with clinical outcome in Fanconi anemia. European Fanconi Anemia Research Group. *Blood*. 2000;96:4064–70. PubMed PMID: 11110674.
- Faivre L, Portnoi MF, Pals G, Stoppa-Lyonnet D, Le Merrer M, Thauvin-Robinet C, Huet F, Mathew CG, Joenje H, Verloes A, Baumann C. Should chromosome breakage studies be performed in patients with VACTERL association? *Am J Med Genet A*. 2005;137:55–8. PubMed PMID: 16015582.
- Fanconi Anaemia/Breast Cancer Consortium. Positional cloning of the Fanconi anaemia group A gene. *Nat Genet*. 1996;14:324–8. PubMed PMID: 8896564.
- Ferrer M, Rodriguez JA, Spierings EA, de Winter JP, Giaccone G, Kruyt FA. Identification of multiple nuclear export sequences in Fanconi anemia group A protein that contribute to CRM1-dependent nuclear export. *Hum Mol Genet*. 2005;14:1271–81. PubMed PMID: 15790592.
- Frohnmayr D, Frohnmayr L, Guinan E, Kennedy T, Larsen K, eds. *Fanconi Anemia: Guidelines for Diagnosis and Management*. 4 ed. Eugene, OR: Fanconi Anemia Research Fund, Inc. Available [online](#). 2014. Accessed 3-6-18.
- Futaki M, Yamashita T, Yagasaki H, Toda T, Yabe M, Kato S, Asano S, Nakahata T. The IVS4 + 4 A to T mutation of the Fanconi anemia gene FANCC is not associated with a severe phenotype in Japanese patients. *Blood*. 2000;95:1493–8. PubMed PMID: 10666230.
- Garbati MR, Hays LE, Keeble W, Yates JE, Rathbun RK, Bagby GC. FANCA and FANCC modulate TLR and p38 MAPK-dependent expression of IL-1 β in macrophages. *Blood*. 2013;122:3197–205. PubMed PMID: 24046015.

- Garcia-Higuera I, Taniguchi T, Ganesan S, Meyn MS, Timmers C, Hejna J, Grompe M, D'Andrea AD. Interaction of the Fanconi anemia proteins and BRCA1 in a common pathway. *Mol Cell*. 2001;7:249. PubMed PMID: 11239454.
- Gillio AP, Verlander PC, Batish SD, Giampietro PF, Auerbach AD. Phenotypic consequences of mutations in the Fanconi anemia FAC gene: an International Fanconi Anemia Registry study. *Blood*. 1997;90:105–10. PubMed PMID: 9207444.
- Giulino L, Guinan EC, Gillio AP, Drachtman RA, Teruya-Feldstein J, Boulad F. Sweet syndrome in patients with Fanconi anaemia: association with extracutaneous manifestations and progression of haematological disease. *Br J Haematol*. 2011;154:278–81. PubMed PMID: 21501135.
- Guardiola P, Socié G, Li X, Ribaud P, Devergie A, Espérou H, Richard P, Traineau R, Janin A, Gluckman E. Acute graft-versus-host disease in patients with Fanconi anemia or acquired aplastic anemia undergoing bone marrow transplantation from HLA-identical sibling donors: risk factors and influence on outcome. *Blood*. 2004;103:73–7. PubMed PMID: 12946993.
- Guo M, Vidhyasagar V, Ding H, Wu Y. Insight into the roles of helicase motif Ia by characterizing Fanconi anemia group J protein (FANCF) patient mutations. *J Biol Chem*. 2014;289:10551–65. PubMed PMID: 24573678.
- Haitjema A, Mol BM, Kooi IE, Massink MP, Jørgensen JA, Rockx DA, Rooimans MA, de Winter JP, Meijers-Heijboer H, Joenje H, Dorsman JC. Coregulation of FANCA and BRCA1 in human cells. *Springerplus*. 2014;3:381. PubMed PMID: 25161863.
- Hira A, Yoshida K, Sato K, Okuno Y, Shiraishi Y, Chiba K, Tanaka H, Miyano S, Shimamoto A, Tahara H, Ito E, Kojima S, Kurumizaka H, Ogawa S, Takata M, Yabe H, Yabe M. Mutations in the gene encoding the E2 conjugating enzyme UBE2T cause Fanconi anemia. *Am J Hum Genet*. 2015;96:1001–7. PubMed PMID: 26046368.
- Hirsch B, Shimamura A, Moreau L, Baldinger S, Hag-alshiekh M, Bostrom B, Sencer S, D'Andrea AD. Association of biallelic BRCA2/FANCD1 mutations with spontaneous chromosomal instability and solid tumors of childhood. *Blood*. 2004;103:2554–9. PubMed PMID: 14670928.
- Ho GP, Margossian S, Taniguchi T, D'Andrea AD. Phosphorylation of FANCD2 on two novel sites is required for mitomycin C resistance. *Mol Cell Biol*. 2006;26:7005–15. PubMed PMID: 16943440.
- Ho TV, Schärer OD. Translesion DNA synthesis polymerases in DNA interstrand crosslink repair. *Environ Mol Mutagen*. 2010;51:552–66. PubMed PMID: 20658647.
- Hosoya Y, Lefor A, Hirashima Y, Nokubi M, Yamaguti T, Jinbu Y, Muroi K, Nakazawa M, Yasuda Y. Successful treatment of esophageal squamous cell carcinoma in a patient with Fanconi anemia. *Jpn J Clin Oncol*. 2010;40:805–10. PubMed PMID: 20410055.
- Houghtaling S, Timmers C, Noll M, Finegold MJ, Jones SN, Meyn MS, Grompe M. Epithelial cancer in Fanconi anemia complementation group D2 (Fancd2) knockout mice. *Genes Dev*. 2003;17:2021–35. PubMed PMID: 12893777.
- Howlett NG, Taniguchi T, Olson S, Cox B, Waisfisz Q, De Die-Smulders C, Persky N, Grompe M, Joenje H, Pals G, Ikeda H, Fox EA, D'Andrea AD. Biallelic inactivation of BRCA2 in Fanconi anemia. *Science*. 2002;297:606–9. PubMed PMID: 12065746.
- Howlett NG, Harney JA, Rego MA, Kolling FW 4th, Glover TW. Functional interaction between the Fanconi Anemia D2 protein and proliferating cell nuclear antigen (PCNA) via a conserved putative PCNA interaction motif. *J Biol Chem*. 2009;284:28935–42. PubMed PMID: 19704162.
- Huang M, D'Andrea AD. A new nuclease member of the FAN club. *Nat Struct Mol Biol*. 2010;17:926–8. PubMed PMID: 20683477.

- Huang M, Kim JM, Shiotani B, Yang K, Zou L, D'Andrea AD. The FANCM/FAAP24 complex is required for the DNA interstrand crosslink-induced checkpoint response. *Mol Cell*. 2010;39:259–68. PubMed PMID: 20670894.
- Huang Y, Leung JW, Lowery M, Matsushita N, Wang Y, Shen X, Huong D, Takata M, Chen J, Li L. Modularized functions of the Fanconi anemia core complex. *Cell Rep*. 2014;7:1849–57. PubMed PMID: 24910428.
- Huang TT, Nijman SM, Mirchandani KD, Galardy PJ, Cohn MA, Haas W, Gygi SP, Ploegh HL, Bernardis R, D'Andrea AD. Regulation of monoubiquitinated PCNA by DUB autocleavage. *Nat Cell Biol*. 2006;8:339–47. PubMed PMID: 16531995.
- Ishiai M, Kitao H, Smogorzewska A, Tomida J, Kinomura A, Uchida E, Saberi A, Kinoshita E, Kinoshita-Kikuta E, Koike T, Tashiro S, Elledge SJ, Takata M. FANCI phosphorylation functions as a molecular switch to turn on the Fanconi anemia pathway. *Nat Struct Mol Biol*. 2008;15:1138–46. PubMed PMID: 18931676.
- Jones S, Hruban RH, Kamiyama M, Borges M, Zhang X, Parsons DW, Lin JC, Palmisano E, Brune K, Jaffee EM, Iacobuzio-Donahue CA, Maitra A, Parmigiani G, Kern SE, Velculescu VE, Kinzler KW, Vogelstein B, Eshleman JR, Goggins M, Klein AP. Exomic sequencing identifies PALB2 as a pancreatic cancer susceptibility gene. *Science*. 2009;324:217. PubMed PMID: 19264984.
- Joo W, Xu G, Persky NS, Smogorzewska A, Rudge DG, Buzovetsky O, Elledge SJ, Pavletich NP. Structure of the FANCI-FANCD2 complex: insights into the Fanconi anemia DNA repair pathway. *Science*. 2011;333:312–6. PubMed PMID: 21764741.
- Kahraman S, Beyazyurek C, Yesilipek MA, Ozturk G, Ertem M, Anak S, Kansoy S, Aksoylar S, Kuşkonmaz B, Oniz H, Slavin S, Karakas Z, Tac HA, Gulum N, Ekmekci GC. Successful haematopoietic stem cell transplantation in 44 children from healthy siblings conceived after preimplantation HLA matching. *Reprod Biomed Online*. 2014;29:340–51. PubMed PMID: 25066893.
- Kee Y, D'Andrea AD. Molecular pathogenesis and clinical management of Fanconi anemia. *J Clin Invest*. 2012;122:3799–806. PubMed PMID: 23114602.
- Kim H, D'Andrea AD. Regulation of DNA cross-link repair by the Fanconi anemia/BRCA pathway. *Genes Dev*. 2012;26:1393–408. PubMed PMID: 22751496.
- Kim Y, Lach FP, Desetty R, Hanenberg H, Auerbach AD, Smogorzewska A. Mutations of the SLX4 gene in Fanconi anemia. *Nat Genet*. 2011;43:142–6. PubMed PMID: 21240275.
- Knies K, Inano S, Ramírez MJ, Ishiai M, Surrallés J, Takata M, Schindler D. Biallelic mutations in the ubiquitin ligase RFWF3 cause Fanconi anemia. *J Clin Invest*. 2017;127:3013–27. PubMed PMID: 28691929.
- Knipscheer P, Räschle M, Smogorzewska A, Enoiu M, Ho TV, Schärer OD, Elledge SJ, Walter JC. The Fanconi anemia pathway promotes replication-dependent DNA interstrand cross-link repair. *Science*. 2009;326:1698–701. PubMed PMID: 19965384.
- Kowal P, Gurtan AM, Stuckert P, D'Andrea AD, Ellenberger T. Structural determinants of human FANCF protein that function in the assembly of a DNA damage signaling complex. *J Biol Chem*. 2007;282:2047–55. PubMed PMID: 17082180.
- Kratz K, Schöpf B, Kaden S, Sendoel A, Eberhard R, Lademann C, Cannavó E, Sartori AA, Hengartner MO, Jiricny J. Deficiency of FANCD2-associated nuclease KIAA1018/FAN1 sensitizes cells to interstrand crosslinking agents. *Cell*. 2010;142:77–88. PubMed PMID: 20603016.
- Kutler DI, Singh B, Satagopan J, Batish SD, Berwick M, Giampietro PF, Hanenberg H, Auerbach AD. A 20-year perspective on the International Fanconi Anemia Registry (IFAR). *Blood*. 2003;101:1249–56. PubMed PMID: 12393516.
- Leung JW, Wang Y, Fong KW, Huen MS, Li L, Chen J. Fanconi anemia (FA) binding protein FAAP20 stabilizes FA complementation group A (FANCA) and participates in interstrand cross-link repair. *Proc Natl Acad Sci U S A*. 2012;109:4491–6. PubMed PMID: 22396592.

- Léveillé F, Blom E, Medhurst AL, Bier P, Laghmani el H, Johnson M, Rooimans MA, Soback A, Waisfisz Q, Arwert F, Patel KJ, Hoatlin ME, Joenje H, de Winter JP. The Fanconi anemia gene product FANCF is a flexible adaptor protein. *J Biol Chem.* 2004;279:39421–30. PubMed PMID: 15262960.
- Léveillé F, Ferrer M, Medhurst AL, Laghmani el H, Rooimans MA, Bier P, Steltenpool J, Titus TA, Postlethwait JH, Hoatlin ME, Joenje H, de Winter JP. The nuclear accumulation of the Fanconi anemia protein FANCE depends on FANCC. *DNA Repair (Amst).* 2006;5:556–65. PubMed PMID: 16513431.
- Levitus M, Waisfisz Q, Godthelp BC, de Vries Y, Hussain S, Wiegant WW, Elghalbzouri-Maghrani E, Steltenpool J, Rooimans MA, Pals G, Arwert F, Mathew CG, Zdzienicka MZ, Hiom K, De Winter JP, Joenje H. The DNA helicase BRIP1 is defective in Fanconi anemia complementation group J. *Nat Genet.* 2005;37:934–5. PubMed PMID: 16116423.
- Levrán O, Attwooll C, Henry RT, Milton KL, Neveling K, Rio P, Batish SD, Kalb R, Velleuer E, Barral S, Ott J, Petrini J, Schindler D, Hanenberg H, Auerbach AD. The BRCA1-interacting helicase BRIP1 is deficient in Fanconi anemia. *Nat Genet.* 2005;37:931–3. PubMed PMID: 16116424.
- Levrán O, Erlich T, Magdalena N, Gregory JJ, Batish SD, Verlander PC, Auerbach AD. Sequence variation in the Fanconi anemia gene FAA. *Proc Natl Acad Sci U S A.* 1997;94:13051–6. PubMed PMID: 9371798.
- Liang CC, Zhan B, Yoshikawa Y, Haas W, Gygi SP, Cohn MA. UHRF1 is a sensor for DNA interstrand crosslinks and recruits FANCD2 to initiate the Fanconi anemia pathway. *Cell Rep.* 2015;10:1947–56. PubMed PMID: 25801034.
- Ling C, Ishiai M, Ali AM, Medhurst AL, Neveling K, Kalb R, Yan Z, Xue Y, Oostra AB, Auerbach AD, Hoatlin ME, Schindler D, Joenje H, de Winter JP, Takata M, Meetei AR, Wang W. FAAP100 is essential for activation of the Fanconi anemia-associated DNA damage response pathway. *EMBO J.* 2007;26:2104–14. PubMed PMID: 17396147.
- Litman R, Peng M, Jin Z, Zhang F, Zhang J, Powell S, Andreassen PR, Cantor SB. BACH1 is critical for homologous recombination and appears to be the Fanconi anemia gene product FANCF. *Cancer Cell.* 2005;8:255–65. PubMed PMID: 16153896.
- Lo Ten Foe JR, Rooimans MA, Bosnoyan-Collins L, Alon N, Wijker M, Parker L, Lightfoot J, Carreau M, Callen DF, Savoia A, Cheng NC, van Berkel CG, Strunk MH, Gille JJ, Pals G, Kruyt FA, Pronk JC, Arwert F, Buchwald M, Joenje H. Expression cloning of a cDNA for the major Fanconi anaemia gene, FAA. *Nat Genet.* 1996;14:320–3. PubMed PMID: 8896563.
- Lossaint G, Larroque M, Ribeyre C, Bec N, Larroque C, Décaillot C, Gari K, Constantinou A. FANCD2 binds MCM proteins and controls replisome function upon activation of S phase checkpoint signaling. *Mol Cell.* 2013;51:678–90. PubMed PMID: 23993743.
- MacKay C, Déclais AC, Lundin C, Agostinho A, Deans AJ, MacArtney TJ, Hofmann K, Gartner A, West SC, Helleday T, Lilley DM, Rouse J. Identification of KIAA1018/FAN1, a DNA repair nuclease recruited to DNA damage by monoubiquitinated FANCD2. *Cell.* 2010;142:65–76. PubMed PMID: 20603015.
- MacMillan ML, DeFor TE, Young JA, Dusenbery KE, Blazar BR, Slungaard A, Zierhut H, Weisdorf DJ, Wagner JE. Alternative donor hematopoietic cell transplantation for Fanconi anemia. *Blood.* 2015;125:3798–804. PubMed PMID: 25824692.
- MacMillan ML, Wagner JE. Haematopoietic cell transplantation for Fanconi anaemia - when and how? *Br J Haematol.* 2010;149:14–21. PubMed PMID: 20136826.
- Machida YJ, Machida Y, Chen Y, Gurtan AM, Kupfer GM, D'Andrea AD, Dutta A. UBE2T is the E2 in the Fanconi anemia pathway and undergoes negative autoregulation. *Mol Cell.* 2006;23:589–96. PubMed PMID: 16916645.
- Magron A, Elowe S, Carreau M. The Fanconi Anemia C Protein Binds to and Regulates Stathmin-1 Phosphorylation. *PLoS One.* 2015;10:e0140612. PubMed PMID: 26466335.

- Marsit CJ, Liu M, Nelson HH, Posner M, Suzuki M, Kelsey KT. Inactivation of the Fanconi anemia/BRCA pathway in lung and oral cancers: implications for treatment and survival. *Oncogene*. 2004;23:1000–4. PubMed PMID: 14647419.
- Masserot C, Peffault de Latour R, Rocha V, Leblanc T, Rigolet A, Pascal F, Janin A, Soulier J, Gluckman E, Socié G. Head and neck squamous cell carcinoma in 13 patients with Fanconi anemia after hematopoietic stem cell transplantation. *Cancer*. 2008;113:3315–22. PubMed PMID: 18831513.
- Matsushita N, Kitao H, Ishiai M, Nagashima N, Hirano S, Okawa K, Ohta T, Yu DS, McHugh PJ, Hickson ID, Venkitaraman AR, Kurumizaka H, Takata M. A FancD2-monoubiquitin fusion reveals hidden functions of Fanconi anemia core complex in DNA repair. *Mol Cell*. 2005;19:841–7. PubMed PMID: 16168378.
- McCauley J, Masand N, McGowan R, Rajagopalan S, Hunter A, Michaud JL, Gibson K, Robertson J, Vaz F, Abbs S, Holden ST. X-linked VACTERL with hydrocephalus syndrome: further delineation of the phenotype caused by FANCB mutations. *Am J Med Genet A*. 2011;155A:2370–80. PubMed PMID: 21910217.
- Meetei AR, de Winter JP, Medhurst AL, Wallisch M, Waisfisz Q, van de Vrugt HJ, Oostra AB, Yan Z, Ling C, Bishop CE, Hoatlin ME, Joenje H, Wang W. A novel ubiquitin ligase is deficient in Fanconi anemia. *Nat Genet*. 2003a;35:165–70. PubMed PMID: 12973351.
- Meetei AR, Levitus M, Xue Y, Medhurst AL, Zwaan M, Ling C, Rooimans MA, Bier P, Hoatlin M, Pals G, de Winter JP, Wang W, Joenje H. X-linked inheritance of Fanconi anemia complementation group B. *Nat Genet*. 2004;36:1219–24. PubMed PMID: 15502827.
- Meetei AR, Medhurst AL, Ling C, Xue Y, Singh TR, Bier P, Steltenpool J, Stone S, Dokal I, Mathew CG, Hoatlin M, Joenje H, de Winter JP, Wang W. A human ortholog of archaeal DNA repair protein Hef is defective in Fanconi anemia complementation group M. *Nat Genet*. 2005;37:958–63. PubMed PMID: 16116422.
- Meetei AR, Sechi S, Wallisch M, Yang D, Young MK, Joenje H, Hoatlin ME, Wang W. A multiprotein nuclear complex connects Fanconi anemia and Bloom syndrome. *Mol Cell Biol*. 2003b;23:3417–26. PubMed PMID: 12724401.
- Mehta PA, Davies SM, Leemhuis T, Myers K, Kernan NA, Prockop SE, Scaradavou A, O'Reilly RJ, Williams DA, Lehmann L, Guinan E, Margolis D, Baker KS, Lane A, Boulad F. Radiation-free, alternative donor HCT for Fanconi anemia patients: results from a prospective multi-institutional study. *Blood*. 2017;129:2308–15. PubMed PMID: 28179273.
- Mehta PA, Harris RE, Davies SM, Kim MO, Mueller R, Lampkin B, Mo J, Myers K, Smolarek TA. Numerical chromosomal changes and risk of development of myelodysplastic syndrome—acute myeloid leukemia in patients with Fanconi anemia. *Cancer Genet Cytogenet*. 2010;203:180–6. PubMed PMID: 21156231.
- Mehta PA, Ileri T, Harris RE, Williams DA, Mo J, Smolarek T, Auerbach AD, Kelly P, Davies SM. Chemotherapy for myeloid malignancy in children with Fanconi anemia. *Pediatr Blood Cancer*. 2007;48:668–72. PubMed PMID: 16609946.
- Meindl A, Hellebrand H, Wiek C, Erven V, Wappenschmidt B, Niederacher D, Freund M, Lichtner P, Hartmann L, Schaal H, Ramser J, Honisch E, Kubisch C, Wichmann HE, Kast K, Deissler H, Engel C, Müller-Myhsok B, Neveling K, Kiechle M, Mathew CG, Schindler D, Schmutzler RK, Hanenberg H. Germline mutations in breast and ovarian cancer pedigrees establish RAD51C as a human cancer susceptibility gene. *Nat Genet*. 2010;42:410–4. PubMed PMID: 20400964.
- Mi J, Qiao F, Wilson JB, High AA, Schroeder MJ, Stukenberg PT, Moss A, Shabanowitz J, Hunt DF, Jones NJ, Kupfer GM. FANCG is phosphorylated at serines 383 and 387 during mitosis. *Mol Cell Biol*. 2004;24:8576–85. PubMed PMID: 15367677.
- Montes de Oca R, Andreassen PR, Margossian SP, Gregory RC, Taniguchi T, Wang X, Houghtaling S, Grompe M, D'Andrea AD. Regulated interaction of the Fanconi anemia protein, FANCD2, with chromatin. *Blood*. 2005;105:1003–9. PubMed PMID: 15454491.

- Morgan NV, Tipping AJ, Joenje H, Mathew CG. High frequency of large intragenic deletions in the Fanconi anemia group A gene. *Am J Hum Genet.* 1999;65:1330–41. PubMed PMID: 10521298.
- Morris LG, Sikora AG, Patel SG, Hayes RB, Ganly I. Second primary cancers after an index head and neck cancer: subsite-specific trends in the era of human papillomavirus-associated oropharyngeal cancer. *J Clin Oncol.* 2011;29:739–46. PubMed PMID: 21189382.
- Moynahan ME, Pierce AJ, Jasin M. BRCA2 is required for homology-directed repair of chromosomal breaks. *Mol Cell.* 2001;7:263–72. PubMed PMID: 11239455.
- Myers K, Davies SM, Harris RE, Spunt SL, Smolarek T, Zimmerman S, McMasters R, Wagner L, Mueller R, Auerbach AD, Mehta PA. The clinical phenotype of children with Fanconi anemia caused by biallelic FANCD1/BRCA2 mutations. *Pediatr Blood Cancer.* 2012;58:462–5. PubMed PMID: 21548014.
- Nabhan SK, Bitencourt MA, Duval M, Abecasis M, Dufour C, Boudjedir K, Rocha V, Socié G, Passweg J, Goi K, Sanders J, Snowden J, Yabe H, Pasquini R, Gluckman E; Aplastic Anaemia Working Party, EBMT. Fertility recovery and pregnancy after allogeneic hematopoietic stem cell transplantation in Fanconi anemia patients. *Haematologica.* 2010;95:1783–7. PubMed PMID: 20494929.
- Nakanishi K, Cavallo F, Perrouault L, Giovannangeli C, Moynahan ME, Barchi M, Brunet E, Jasin M. Homology-directed Fanconi anemia pathway cross-link repair is dependent on DNA replication. *Nat Struct Mol Biol.* 2011;18:500–3. PubMed PMID: 21423196.
- Nakanishi K, Taniguchi T, Ranganathan V, New HV, Moreau LA, Stotsky M, Mathew CG, Kastan MB, Weaver DT, D'Andrea AD. Interaction of FANCD2 and NBS1 in the DNA damage response. *Nat Cell Biol.* 2002;4:913–20. PubMed PMID: 12447395.
- Narayan G, Arias-Pulido H, Nandula SV, Basso K, Sugirtharaj DD, Vargas H, Mansukhani M, Vilella J, Meyer L, Schneider A, Gissmann L, Dürst M, Pothuri B, Murty VV. Promoter hypermethylation of FANCF: disruption of Fanconi Anemia-BRCA pathway in cervical cancer. *Cancer Res.* 2004;64:2994–7. PubMed PMID: 15126331.
- Neitzel H, Kühl JS, Gerlach A, Ebell W, Tönnies H. Clonal chromosomal aberrations in bone marrow cells of Fanconi anemia patients: results and implications. In: Schindler D, Hoehn H, eds. *Fanconi Anemia. a Paradigmatic Disease for the Understanding of Cancer and Aging.* Monographs in Human Genetics. Vol 15. Basel: Karger. 2007:79-94
- Neveling K, Endt D, Hoehn H, Schindler D. Genotype–phenotype correlations in Fanconi anemia. *Mutat Res.* 2009;668:73–91. PubMed PMID: 19464302.
- Nicchia E, Benedicenti F, De Rocco D, Greco C, Bottega R, Inzana F, Faleschini M, Bonin S, Cappelli E, Moggi M, Stanzial F, Svahn J, Dufour C, Savoia A. Clinical aspects of Fanconi anemia individuals with the same mutation of FANCF identified by next generation sequencing. *Birth Defects Res A Clin Mol Teratol.* 2015;103:1003–10. PubMed PMID: 26033879.
- Nijman SM, Huang T'T, Dirac AM, Brummelkamp TR, Kerkhoven RM, D'Andrea AD, Bernards R. The deubiquitinating enzyme USP1 regulates the Fanconi anemia pathway. *Mol Cell.* 2005;17:331–9. PubMed PMID: 15694335.
- Nookala RK, Hussain S, Pellegrini L. Insights into Fanconi Anaemia from the structure of human FANCE. *Nucleic Acids Res.* 2007;35:1638–48. PubMed PMID: 17308347.
- Oda T, Hayano T, Miyaso H, Takahashi N, Yamashita T. Hsp90 regulates the Fanconi anemia DNA damage response pathway. *Blood.* 2007;109:5016–26. PubMed PMID: 17327415.
- Oestergaard VH, Langevin F, Kuiken HJ, Pace P, Niedzwiedz W, Simpson LJ, Ohzeki M, Takata M, Sale JE, Patel KJ. Deubiquitination of FANCD2 is required for DNA crosslink repair. *Mol Cell.* 2007;28:798–809. PubMed PMID: 18082605.

- Olopade OI, Wei M. FANCF methylation contributes to chemoselectivity in ovarian cancer. *Cancer Cell*. 2003;3:417–20. PubMed PMID: 12781358.
- Osorio A, Bogliolo M, Fernández V, Barroso A, de la Hoya M, Caldés T, Lasa A, Ramón y Cajal T, Santamariña M, Vega A, Quiles F, Lázaro C, Díez O, Fernández D, González-Sarmiento R, Durán M, Piqueras JF, Marín M, Pujol R, Surrallés J, Benítez J. Evaluation of rare variants in the new fanconi anemia gene ERCC4 (FANCC) as familial breast/ovarian cancer susceptibility alleles. *Hum Mutat*. 2013;34:1615–8. PubMed PMID: 24027083.
- Pace P, Johnson M, Tan WM, Mosedale G, Sng C, Hoatlin M, de Winter J, Joenje H, Gergely F, Patel KJ. FANCE: the link between Fanconi anaemia complex assembly and activity. *EMBO J*. 2002;21:3414–23. PubMed PMID: 12093742.
- Pang Q, Fagerlie S, Christianson TA, Keeble W, Faulkner G, Diaz J, Rathbun RK, Bagby GC. The Fanconi anemia protein FANCC binds to and facilitates the activation of STAT1 by gamma interferon and hematopoietic growth factors. *Mol Cell Biol*. 2000;20:4724–35. PubMed PMID: 10848598.
- Park JY, Virts EL, Jankowska A, Wiek C, Othman M, Chakraborty SC, Vance GH, Alkuraya FS, Hanenberg H, Andreassen PR. Complementation of hypersensitivity to DNA interstrand crosslinking agents demonstrates that XRCC2 is a Fanconi anaemia gene. *J Med Genet*. 2016;53:672–80. PubMed PMID: 27208205.
- Parmar K, Kim J, Sykes SM, Shimamura A, Stuckert P, Zhu K, Hamilton A, Deloach MK, Kutok JL, Akashi K, Gilliland DG, D'andrea A. Hematopoietic stem cell defects in mice with deficiency of Fancd2 or Usp1. *Stem Cells*. 2010;28:1186–95. PubMed PMID: 20506303.
- Pichierri P, Rosselli F. The DNA crosslink-induced S-phase checkpoint depends on ATR-CHK1 and ATR-NBS1-FANCD2 pathways. *EMBO J*. 2004;23:1178–87. PubMed PMID: 14988723.
- Polito D, Cukras S, Wang X, Spence P, Moreau L, D'Andrea AD, Kee Y. The carboxyl terminus of FANCE recruits FANCD2 to the Fanconi anemia (FA) E3 ligase complex to promote the FA DNA repair pathway. *J Biol Chem*. 2014;289:7003–10. PubMed PMID: 24451376.
- Qiao F, Mi J, Wilson JB, Zhi G, Bucheimer NR, Jones NJ, Kupfer GM. Phosphorylation of Fanconi anemia (FA) complementation group G protein, FANCG, at serine 7 is important for function of the FA pathway. *J Biol Chem*. 2004;279:46035–45. PubMed PMID: 15299017.
- Rafnar T, Gudbjartsson DF, Sulem P, Jonasdottir A, Sigurdsson A, Jonasdottir A, Besenbacher S, Lundin P, Stacey SN, Gudmundsson J, Magnusson OT, le Roux L, Orlygsdottir G, Helgadottir HT, Johannsdottir H, Gylfason A, Tryggvadottir L, Jonasson JG, de Juan A, Ortega E, Ramon-Cajal JM, García-Prats MD, Mayordomo C, Panadero A, Rivera F, Aben KK, van Altena AM, Massuger LF, Aavikko M, Kujala PM, Staff S, Aaltonen LA, Olafsdottir K, Bjornsson J, Kong A, Salvarsdottir A, Saemundsson H, Olafsson K, Benediktsdottir KR, Gulcher J, Masson G, Kiemeny LA, Mayordomo JI, Thorsteinsdottir U, Stefansson K. Mutations in BRIP1 confer high risk of ovarian cancer. *Nat Genet*. 2011;43:1104–7. PubMed PMID: 21964575.
- Rahman N, Seal S, Thompson D, Kelly P, Renwick A, Elliott A, Reid S, Spanova K, Barfoot R, Chagtai T, Jayatilake H, McGuffog L, Hanks S, Evans DG, Eccles D. Breast Cancer Susceptibility Collaboration (UK), Easton DF, Stratton MR. PALB2, which encodes a BRCA2-interacting protein, is a breast cancer susceptibility gene. *Nat Genet*. 2007;39:165–7. PubMed PMID: 17200668.
- Rajendra E, Oestergaard VH, Langevin F, Wang M, Dornan GL, Patel KJ, Passmore LA. The genetic and biochemical basis of FANCD2 monoubiquitination. *Mol Cell*. 2014;54:858–69. PubMed PMID: 24905007.
- Reid S, Schindler D, Hanenberg H, Barker K, Hanks S, Kalb R, Neveling K, Kelly P, Seal S, Freund M, Wurm M, Batish SD, Lach FP, Yetgin S, Neitzel H, Ariffin H, Tischkowitz M, Mathew CG, Auerbach AD, Rahman N. Biallelic mutations in PALB2 cause Fanconi anemia subtype FA-N and predispose to childhood cancer. *Nat Genet*. 2007;39:162–4. PubMed PMID: 17200671.

- Rickman KA, Lach FP, Abhyankar A, Donovan FX, Sanborn EM, Kennedy JA, Sougnez C, Gabriel SB, Elemento O, Chandrasekharappa SC, Schindler D, Auerbach AD, Smogorzewska A. Deficiency of UBE2T, the E2 ubiquitin ligase necessary for FANCD2 and FANCI ubiquitination, causes FA-T subtype of Fanconi anemia. *Cell Rep.* 2015;12:35–41. PubMed PMID: 26119737.
- Rose SR, Kim MO, Korbee L, Wilson KA, Douglas Ris M, Eyal O, Sherafat-Kazemzadeh R, Bollepalli S, Harris R, Jeng MR, Williams DA, Smith FO. Oxandrolone for the treatment of bone marrow failure in Fanconi anemia. *Pediatr Blood Cancer.* 2014;61:11–9. PubMed PMID: 24019220.
- Rosenberg PS, Alter BP, Ebell W. Cancer risks in Fanconi anemia: findings from the German Fanconi Anemia Registry. *Haematologica.* 2008;93:511–7. PubMed PMID: 18322251.
- Rosenberg PS, Socié G, Alter BP, Gluckman E. Risk of head and neck squamous cell cancer and death in patients with Fanconi anemia who did and did not receive transplants. *Blood.* 2005;105:67–73. PubMed PMID: 15331448.
- Rosenberg PS, Tamary H, Alter BP. How high are carrier frequencies of rare recessive syndromes? Contemporary estimates for Fanconi Anemia in the United States and Israel. *Am J Med Genet A.* 2011;155A:1877–83. PubMed PMID: 21739583.
- Sarkies P, Murat P, Phillips LG, Patel KJ, Balasubramanian S, Sale JE. FANCI coordinates two pathways that maintain epigenetic stability at G-quadruplex DNA. *Nucleic Acids Res.* 2012;40:1485–98. PubMed PMID: 22021381.
- Sato K, Toda K, Ishiai M, Takata M, Kurumizaka H. DNA robustly stimulates FANCD2 monoubiquitylation in the complex with FANCI. *Nucleic Acids Res.* 2012;40:4553–61. PubMed PMID: 22287633.
- Sawyer SL, Tian L, Kähkönen M, Schwartzentruber J, Kircher M, et al. Biallelic mutations in BRCA1 cause a new Fanconi anemia subtype. *Cancer Discov.* 2015;5:135–42. PubMed PMID: 25472942.
- Scheckenbach K, Morgan M, Filger-Brillinger J, Sandmann M, Strimling B, Scheurlen W, Schindler D, Gobel U, Hanenberg H. Treatment of the bone marrow failure in Fanconi anemia patients with danazol. *Blood Cells Mol Dis.* 2012;48:128–131. PubMed PMID: 22178060.
- Seal S, Thompson D, Renwick A, Elliott A, Kelly P, Barfoot R, Chagtai T, Jayatilake H, Ahmed M, Spanova K, North B, McGuffog L, Evans DG, Eccles D, et al. Truncating mutations in the Fanconi anemia J gene BRIP1 are low-penetrance breast cancer susceptibility alleles. *Nat Genet.* 2006;38:1239–41. PubMed PMID: 17033622.
- Shimamura A, Alter BP. Pathophysiology and management of inherited bone marrow failure syndromes. *Blood Rev.* 2010;24:101–22. PubMed PMID: 20417588.
- Singh TR, Bakker ST, Agarwal S, Jansen M, Grassman E, Godthelp BC, Ali AM, Du CH, Rooimans MA, Fan Q, Wahengbam K, Steltenpool J, Andreassen PR, Williams DA, Joenje H, de Winter JP, Meetei AR. Impaired FANCD2 monoubiquitination and hypersensitivity to camptothecin uniquely characterize Fanconi anemia complementation group M. *Blood.* 2009;114:174–80. PubMed PMID: 19423727.
- Smogorzewska A, Desetty R, Saito TT, Schlabach M, Lach FP, Sowa ME, Clark AB, Kunkel TA, Harper JW, Colaiácovo MP, Elledge SJ. A genetic screen identifies FAN1, a Fanconi anemia-associated nuclease necessary for DNA interstrand crosslink repair. *Mol Cell.* 2010;39:36–47. PubMed PMID: 20603073.
- Smogorzewska A, Matsuoka S, Vinciguerra P, McDonald ER 3rd, Hurov KE, Luo J, Ballif BA, Gygi SP, Hofmann K, D'Andrea AD, Elledge SJ. Identification of the FANCI protein, a monoubiquitinated FANCD2 paralog required for DNA repair. *Cell.* 2007;129:289–301. PubMed PMID: 17412408.
- Sommers JA, Banerjee T, Hinds T, Wan B, Wold MS, Lei M, Brosh RM Jr. Novel function of the Fanconi anemia group J or RECQ1 helicase to disrupt protein-DNA complexes in a replication protein A-stimulated manner. *J Biol Chem.* 2014;289:19928–41. PubMed PMID: 24895130.

- Song IY, Palle K, Gurkar A, Tateishi S, Kupfer GM, Vaziri C. Rad18-mediated translesion synthesis of bulky DNA adducts is coupled to activation of the Fanconi anemia DNA repair pathway. *J Biol Chem*. 2010;285:31525–36. PubMed PMID: 20675655.
- Spanier G, Pohl F, Giese T, Meier JK, Koelbl O, Reichert TE. Fatal course of tonsillar squamous cell carcinoma associated with Fanconi anaemia: a mini review. *J Craniomaxillofac Surg*. 2012;40:510–5. PubMed PMID: 21925890.
- Stoepker C, Hain K, Schuster B, Hilhorst-Hofstee Y, Rooimans MA, Steltenpool J, Oostra AB, Eirich K, Korthof ET, Nieuwint AW, Jaspers NG, Bettecken T, Joenje H, Schindler D, Rouse J, de Winter JP. SLX4, a coordinator of structure-specific endonucleases, is mutated in a new Fanconi anemia subtype. *Nat Genet*. 2011;43:138–41. PubMed PMID: 21240277.
- Talbot A, Peffault de Latour R, Raffoux E, Buchbinder N, Vigouroux S, Milpied N, Leblanc T, Soulier J, Michallet M, Socié G. Sequential treatment for allogeneic hematopoietic stem cell transplantation in Fanconi anemia with acute myeloid leukemia. *Haematologica*. 2014;99:e199–200. PubMed PMID: 25085358.
- Tamary H, Nishri D, Yacobovich J, Zilber R, Dgany O, Krasnov T, Aviner S, Stepensky P, Ravel-Vilk S, Bitan M, Kaplinsky C, Ben Barak A, Elhasid R, Kapelusnik J, Koren A, Levin C, Attias D, Laor R, Yaniv I, Rosenberg PS, Alter BP. Frequency and natural history of inherited bone marrow failure syndromes: the Israeli Inherited Bone Marrow Failure Registry. *Haematologica*. 2010;95:1300–7. PubMed PMID: 20435624.
- Tan IB, Cutcutache I, Zang ZJ, Iqbal J, Yap SF, Hwang W, Lim WT, Teh BT, Rozen S, Tan EH, Tan P. Fanconi's anemia in adulthood: chemoradiation-induced bone marrow failure and a novel FANCA mutation identified by targeted deep sequencing. *J Clin Oncol*. 2011;29:e591–4. PubMed PMID: 21519011.
- Taniguchi T, D'Andrea AD. Molecular pathogenesis of Fanconi anemia: recent progress. *Blood*. 2006;107:4223–33. PubMed PMID: 16493006.
- Taniguchi T, Garcia-Higuera I, Andreassen PR, Gregory RC, Grompe M, D'Andrea AD. S-phase-specific interaction of the Fanconi anemia protein, FANCD2, with BRCA1 and RAD51. *Blood*. 2002a;100:2414–20. PubMed PMID: 12239151.
- Taniguchi T, Garcia-Higuera I, Xu B, Andreassen PR, Gregory RC, Kim ST, Lane WS, Kastan MB, D'Andrea AD. Convergence of the Fanconi anemia and ataxia-telangiectasia signaling pathways. *Cell*. 2002b;109:459–72. PubMed PMID: 12086603.
- Taniguchi T, Tischkowitz M, Ameziane N, Hodgson SV, Mathew CG, Joenje H, Mok SC, D'Andrea AD. Disruption of the Fanconi anemia-BRCA pathway in cisplatin-sensitive ovarian tumors. *Nat Med*. 2003;9:568–74. PubMed PMID: 12692539.
- Tischkowitz M, Xia B, Sabbaghian N, Reis-Filho JS, Hamel N, Li G, van Beers EH, Li L, Khalil T, Quenneville LA, Omeroglu A, Poll A, Lepage P, Wong N, Nederlof PM, Ashworth A, Tonin PN, Narod SA, Livingston DM, Foulkes WD. Analysis of PALB2/FANCN-associated breast cancer families. *Proc Natl Acad Sci U S A*. 2007;104:6788–93. PubMed PMID: 17420451.
- Tolar J, Adair JE, Antoniou M, Bartholomae CC, Becker PS, Blazar BR, Bueren J, Carroll T, Cavazzana-Calvo M, Clapp DW, Dagleish R, Galy A, Gaspar HB, Hanenberg H, Von Kalle C, Kiem HP, Lindeman D, Naldini L, Navarro S, Renella R, Rio P, Sevilla J, Schmidt M, Verhoeven E, Wagner JE, Williams DA, Thrasher AJ. Stem cell gene therapy for fanconi anemia: report from the 1st international Fanconi anemia gene therapy working group meeting. *Mol Ther*. 2011;19:1193–8. PubMed PMID: 21540837.
- Turner N, Tutt A, Ashworth A. Hallmarks of 'BRCAness' in sporadic cancers. *Nat Rev Cancer*. 2004;4:814–9. PubMed PMID: 15510162.
- Umaña LA, Magoulas P, Bi W, Bacino CA. A male newborn with VACTERL association and Fanconi anemia with a FANCB deletion detected by array comparative genomic hybridization (aCGH). *Am J Med Genet A*. 2011;155A:3071–4. PubMed PMID: 22052692.

- van der Heijden MS, Yeo CJ, Hruban RH, Kern SE. Fanconi anemia gene mutations in young-onset pancreatic cancer. *Cancer Res.* 2003;63:2585–8. PubMed PMID: 12750283.
- Vandenberg CJ, Gergely F, Ong CY, Pace P, Mallery DL, Hiom K, Patel KJ. BRCA1-independent ubiquitination of FANCD2. *Mol Cell.* 2003;12:247–54. PubMed PMID: 12887909.
- Vaz F, Hanenberg H, Schuster B, Barker K, Wiek C, Erven V, Neveling K, Endt D, Kesterton I, Autore F, Fraternali F, Freund M, Hartmann L, Grimwade D, Roberts RG, Schaal H, Mohammed S, Rahman N, Schindler D, Mathew CG. Mutation of the RAD51C gene in a Fanconi anemia-like disorder. *Nat Genet.* 2010;42:406–9. PubMed PMID: 20400963.
- Vetro A, Iascone M, Limongelli I, Ameziane N, Gana S, Della Mina E, Giussani U, Ciccone R, Forlino A, Pezzoli L, Rooimans MA, van Essen AJ, Messa J, Rizzuti T, Bianchi P, Dorsman J, de Winter JP, Lalatta F, Zuffardi O. Loss-of-function FANCL mutations associate with severe Fanconi anemia overlapping the VACTERL association. *Hum Mutat.* 2015;36:562–8. PubMed PMID: 25754594.
- Virts EL, Jankowska A, Mackay C, Glaas MF, Wiek C, Kelich SL, Lottmann N, Kennedy FM, Marchal C, Lehnert E, Scharf RE, Dufour C, Lanciotti M, Farruggia P, Santoro A, Savasan S, Scheckenbach K, Schipper J, Wagenmann M, Lewis T, Leffak M, Farlow JL, Foroud TM, Honisch E, Niederacher D, Chakraborty SC, Vance GH, Pruss D, Timms KM, Lanchbury JS, Alpi AF, Hanenberg H. AluY-mediated germline deletion, duplication and somatic stem cell reversion in UBE2T defines a new subtype of Fanconi anemia. *Hum Mol Genet.* 2015;24:5093–108. PubMed PMID: 26085575.
- Wagner JE, Tolar J, Levran O, Scholl T, Deffenbaugh A, Satagopan J, Ben-Porat L, Mah K, Batish SD, Kutler DI, MacMillan ML, Hanenberg H, Auerbach AD. Germline mutations in BRCA2: shared genetic susceptibility to breast cancer, early onset leukemia, and Fanconi anemia. *Blood.* 2004;103:3226–9. PubMed PMID: 15070707.
- Wang AT, Smogorzewska A. SnapShot: Fanconi anemia and associated proteins. *Cell.* 2015;160:354–354.e1. PubMed PMID: 25594185.
- Wang X, Andreassen PR, D'Andrea AD. Functional interaction of monoubiquitinated FANCD2 and BRCA2/FANCD1 in chromatin. *Mol Cell Biol.* 2004;24:5850–62. PubMed PMID: 15199141.
- Wang X, Kennedy RD, Ray K, Stuckert P, Ellenberger T, D'Andrea AD. Chk1-mediated phosphorylation of FANCE is required for the Fanconi anemia/BRCA pathway. *Mol Cell Biol.* 2007;27:3098–108. PubMed PMID: 17296736.
- Wang Z, Li M, Lu S, Zhang Y, Wang H. Promoter hypermethylation of FANCF plays an important role in the occurrence of ovarian cancer through disrupting Fanconi anemia-BRCA pathway. *Cancer Biol Ther.* 2006;5:256–60. PubMed PMID: 16418574.
- Wang AT, Kim T, Wagner JE, Conti BA, Lach FP, Huang AL, Molina H, Sanborn EM, Zierhut H, Cornes BK, Abhyankar A, Sougnez C, Gabriel SB, Auerbach AD, Kowalczykowski SC, Smogorzewska A. A Dominant Mutation in Human RAD51 Reveals Its Function in DNA Interstrand Crosslink Repair Independent of Homologous Recombination. *Mol Cell.* 2015;59:478–90. PubMed PMID: 26253028.
- Whitney MA, Saito H, Jakobs PM, Gibson RA, Moses RE, Grompe M. A common mutation in the FACC gene causes Fanconi anaemia in Ashkenazi Jews. *Nat Genet.* 1993;4:202–5. PubMed PMID: 8348157.
- Wijker M, Morgan NV, Herterich S, van Berkel CG, Tipping AJ, Gross HJ, Gille JJ, Pals G, Savino M, Altay C, Mohan S, Dokal I, Cavenagh J, Marsh J, van Weel M, Ortega JJ, Schuler D, Samochatova E, Karwacki M, Bekassy AN, Abecasis M, Ebell W, Kwee ML, de Ravel T, Mathew CG. Heterogeneous spectrum of mutations in the Fanconi anaemia group A gene. *Eur J Hum Genet.* 1999;7:52–9. PubMed PMID: 10094191.
- Williams SA, Longerich S, Sung P, Vaziri C, Kupfer GM. The E3 ubiquitin ligase RAD18 regulates ubiquitylation and chromatin loading of FANCD2 and FANCI. *Blood.* 2011a;117:5078–87. PubMed PMID: 21355096.

- Williams SA, Wilson JB, Clark AP, Mitson-Salazar A, Tomashevski A, Ananth S, Glazer PM, Semmes OJ, Bale AE, Jones NJ, Kupfer GM. Functional and physical interaction between the mismatch repair and FA-BRCA pathways. *Hum Mol Genet.* 2011b;20:4395–410. PubMed PMID: 21865299.
- Wilson JB, Blom E, Cunningham R, Xiao Y, Kupfer GM, Jones NJ. Several tetratricopeptide repeat (TPR) motifs of FANCG are required for assembly of the BRCA2/D1-D2-G-X3 complex, FANCD2 monoubiquitylation and phleomycin resistance. *Mutat Res.* 2010;689:12–20. PubMed PMID: 20450923.
- Wong JC, Alon N, McKerlie C, Huang JR, Meyn MS, Buchwald M. Targeted disruption of exons 1 to 6 of the Fanconi Anemia group A gene leads to growth retardation, strain-specific microphthalmia, meiotic defects and primordial germ cell hypoplasia. *Hum Mol Genet.* 2003;12:2063–76. PubMed PMID: 12913077.
- Wu Y, Brosh RM Jr. FANCI helicase operates in the Fanconi Anemia DNA repair pathway and the response to replicational stress. *Curr Mol Med.* 2009;9:470–82. PubMed PMID: 19519404.
- Wu Y, Sommers JA, Loiland JA, Kitao H, Kuper J, Kisker C, Brosh RM Jr. The Q motif of Fanconi anemia group J protein (FANCI) DNA helicase regulates its dimerization, DNA binding, and DNA repair function. *J Biol Chem.* 2012;287:21699–716. PubMed PMID: 22582397.
- Xia B, Dorsman JC, Ameziane N, de Vries Y, Rooimans MA, Sheng Q, Pals G, Errami A, Gluckman E, Llera J, Wang W, Livingston DM, Joenje H, de Winter JP. Fanconi anemia is associated with a defect in the BRCA2 partner PALB2. *Nat Genet.* 2007;39:159–61. PubMed PMID: 17200672.
- Xia B, Sheng Q, Nakanishi K, Ohashi A, Wu J, Christ N, Liu X, Jasin M, Couch FJ, Livingston DM. Control of BRCA2 cellular and clinical functions by a nuclear partner, PALB2. *Mol Cell.* 2006;22:719–29. PubMed PMID: 16793542.
- Xie J, Peng M, Guillemette S, Quan S, Maniatis S, Wu Y, Venkatesh A, Shaffer SA, Brosh RM Jr, Cantor SB. FANCI/BACH1 acetylation at lysine 1249 regulates the DNA damage response. *PLoS Genet.* 2012;8:e1002786. PubMed PMID: 22792074.
- Yamashita T, Barber DL, Zhu Y, Wu N, D'Andrea AD. The Fanconi anemia polypeptide FACC is localized to the cytoplasm. *Proc Natl Acad Sci U S A.* 1994;91:6712–6. PubMed PMID: 7517562.
- Yamashita T, Wu N, Kupfer G, Corless C, Joenje H, Grompe M, D'Andrea AD. The Clinical variability of Fanconi Anemia (Type C) results from expression of an amino terminal truncated FAC polypeptide with partial activity. *Blood.* 1996;87:4424. PubMed PMID: 8639804.
- Yoshikiyo K, Kratz K, Hirota K, Nishihara K, Takata M, Kurumizaka H, Horimoto S, Takeda S, Jiricny J. KIAA1018/FAN1 nuclease protects cells against genomic instability induced by interstrand cross-linking agents. *Proc Natl Acad Sci U S A.* 2010;107:21553–7. PubMed PMID: 21115814.
- Zhang F, Fan Q, Ren K, Auerbach AD, Andreassen PR. FANCI/BRIP1 recruitment and regulation of FANCD2 in DNA damage responses. *Chromosoma.* 2010;119:637–49. PubMed PMID: 20676667.
- Zhi G, Wilson JB, Chen X, Krause DS, Xiao Y, Jones NJ, Kupfer GM. Fanconi anemia complementation group FANCD2 protein serine 331 phosphorylation is important for Fanconi anemia pathway function and BRCA2 interaction. *Cancer Res.* 2009;69:8775–83. PubMed PMID: 19861535.
- Zhu B, Yan K, Li L, Lin M, Zhang S, He Q, Zheng D, Yang H, Shao G. K63-linked ubiquitination of FANCG is required for its association with the Rap80-BRCA1 complex to modulate homologous recombination repair of DNA interstrand crosslinks. *Oncogene.* 2015a;34:2867–78. PubMed PMID: 25132264.
- Zhu J, Su F, Mukherjee S, Mori E, Hu B, Asaithamby A. FANCD2 influences replication fork processes and genome stability in response to clustered DSBs. *Cell Cycle.* 2015b;14:1809–22. PubMed PMID: 26083937.
- Zou J, Tian F, Li J, Pickner W, Long M, Rezvani K, Wang H, Zhang D. FANCI regulates interstrand crosslinker induced centrosome amplification through the activation of polo-like kinase 1. *Biol Open.* 2013;2:1022–31. PubMed PMID: 24167712.

Zou J, Zhang D, Qin G, Chen X, Wang H, Zhang D. BRCA1 and FancJ cooperatively promote interstrand crosslinker induced centrosome amplification through the activation of polo-like kinase 1. *Cell Cycle*. 2014;13:3685–97. PubMed PMID: 25483079.

Chapter Notes

Author Notes

Comprehensive Center for Fanconi Anemia

Dana Farber Cancer Institute
Mayer 676
44 Binney Street
Boston, MA 02115
Phone: 617-632-6302

Author History

Blanche P Alter, MD, MPH, FAAP; National Cancer Institute (2011-2016)
Alan D'Andrea, MD; Dana Farber Cancer Institute (2002-2007)
Gary Kupfer, MD; Yale University School of Medicine (2011-2016)
Parinda A Mehta, MD (2016-present)
Lisa Moreau, MS; Dana Farber Cancer Institute (2002-2007)
Akiko Shimamura, MD, PhD; Dana Farber Cancer Institute (2002-2007)
Toshiyasu Taniguchi, MD, PhD; Fred Hutchinson Cancer Research Center (2007-2011)
Jakub Tolar, MD, PhD (2016-present)

Revision History

- 8 March 2018 (aa, pm) Revision: addition of one gene: *RFWD3*
- 23 February 2017 (aa, pm) Revision: addition of two genes: *MAD2L2* and *XRCC2*; edits to Prevention of Secondary Complications
- 22 September 2016 (sw) Comprehensive update posted live
- 7 February 2013 (cd) Revision: deletion/duplication analysis available clinically for *FANCC*
- 6 September 2012 (cd) Revision: sequence analysis for mutations in *RAD51C* and *SLX4* available clinically
- 3 November 2011 (cd) Revision: deletion/duplication analysis available clinically for *PALB2* deletions
- 10 February 2011 (me) Comprehensive update posted live
- 27 March 2008 (cd) Revision: sequence analysis and prenatal testing available clinically for *FANCB*-, *FANCE*-, *FANCF*- and *FANCI*-related Fanconi anemia
- 29 January 2008 (cd) Revision: sequence analysis of entire coding region of *FANCG* and prenatal testing available
- 7 November 2007 (cd) Revision: molecular genetic testing and prenatal diagnosis no longer available on a clinical basis for *FANCF* and *FANCG*
- 22 June 2007 (me) Comprehensive update posted live
- 1 March 2006 (cd) Revision: *FANCB* mutations: X-linked inheritance
- 3 January 2006 (as) Revision: deletion/duplication testing clinically available
- 13 September 2004 (me) Comprehensive update posted live
- 14 February 2002 (me) Review posted live
- 31 May 2001 (as) Original submission

License

GeneReviews® chapters are owned by the University of Washington. Permission is hereby granted to reproduce, distribute, and translate copies of content materials for noncommercial research purposes only, provided that (i) credit for source (<http://www.genereviews.org/>) and copyright (© 1993-2020 University of Washington) are included with each copy; (ii) a link to the original material is provided whenever the material is published elsewhere on the Web; and (iii) reproducers, distributors, and/or translators comply with the [GeneReviews® Copyright Notice and Usage Disclaimer](#). No further modifications are allowed. For clarity, excerpts of GeneReviews chapters for use in lab reports and clinic notes are a permitted use.

For more information, see the [GeneReviews® Copyright Notice and Usage Disclaimer](#).

For questions regarding permissions or whether a specified use is allowed, contact: admasst@uw.edu.