




Exercise and sports participation in patients with thoracic aortic disease: a review

Carlijn G. E. Thijssen, Lidia R. Bons, Arjen L. Gökalp, Roland R. J. Van Kimmenade, Mostafa M. Mokhles, Antonio Pelliccia, Johanna J. M. Takkenberg & Jolien W. Roos-Hesselink


To cite this article: Carlijn G. E. Thijssen, Lidia R. Bons, Arjen L. Gökalp, Roland R. J. Van Kimmenade, Mostafa M. Mokhles, Antonio Pelliccia, Johanna J. M. Takkenberg & Jolien W. Roos-Hesselink (2019) Exercise and sports participation in patients with thoracic aortic disease: a review, Expert Review of Cardiovascular Therapy, 17:4, 251-266, DOI: [10.1080/14779072.2019.1585807](https://doi.org/10.1080/14779072.2019.1585807)


To link to this article: <https://doi.org/10.1080/14779072.2019.1585807>

 View supplementary material [↗](#)


 Accepted author version posted online: 19 Mar 2019.
Published online: 21 Mar 2019.

 Submit your article to this journal [↗](#)

 Article views: 439

 View related articles [↗](#)

 View Crossmark data [↗](#)

 Citing articles: 9 View citing articles [↗](#)

REVIEW



Exercise and sports participation in patients with thoracic aortic disease: a review

Carlijn G. E. Thijssen ^a, Lidia R. Bons^a, Arjen L. Gökalp^b, Roland R. J. Van Kimmenade^c, Mostafa M. Mokhles^b, Antonio Pelliccia^d, Johanna J. M. Takkenberg^b and Jolien W. Roos-Hesselink ^a

^aDepartment of Cardiology, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands; ^bDepartment of Cardiothoracic Surgery, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands; ^cDepartment of Cardiology, Radboud University Medical Center, Nijmegen, The Netherlands; ^dDepartment of Cardiology, Institute of Sports Medicine & Science, Rome, Italy

ABSTRACT

Introduction: Current guidelines recommend patients with thoracic aortic disease (TAD) including inherited aortopathies to avoid heavy exercise. However, evidence supporting the negative advice on exercise is scarce. We aimed to provide an up-to-date systematic review of the available evidence on risks and benefits of exercise and sports participation in TAD patients.

Areas covered: A systematic search was performed in Medline, Embase and Web of Science: thoracic aortic aneurysm or thoracic aortic dissection or inheritable aortopathies including Marfan Syndrome (MFS), Loeys-Dietz syndrome, Turner Syndrome, Ehlers-Danlos syndrome, bicuspid aortic valve (BAV) and sports, exercise or athletes. The resulting 1,652 manuscripts were reviewed by two independent observers. Eventually, 26 studies and 12 case-reports were included, reporting on thoracic aortic dimensions in athletes, exercise related acute aortic dissections, and exercise in BAV and MFS patients.

Expert opinion: Blood pressure elevation during exercise may be associated with an increased risk of acute aortic dissection; however, no controlled trials have longitudinally evaluated the effect of exercise on survival or the risk of aortic dissection in TAD patients. Mouse-model studies suggest beneficial effects of exercise in the setting of a dilated aorta in MFS. There is a clear need for prospective research in this field.

ARTICLE HISTORY

Received 9 November 2018
Accepted 19 February 2019

KEYWORDS

Athletes; bicuspid aortic valve; exercise; heritable thoracic aortic aneurysms and dissection (HTAAD); inheritable aortopathies; Marfan Syndrome; sports participation; thoracic aortic dilatation; thoracic aortic dissection

1. Introduction

The incidence of thoracic aortic disease (TAD) such as thoracic aortic aneurysms and dissections is estimated to be 9.1–16.3/100,000 per year [1]. However, thoracic aortic aneurysms are mostly asymptomatic and its prevalence is probably underestimated. About 20% of patients with thoracic aortic dilatation have a positive family history of aortic disease [2], which can be an expression of an underlying disorder such as bicuspid aortic valve (BAV) or connective tissue disorders, such as Marfan Syndrome (MFS), Loeys-Dietz syndrome, Ehlers-Danlos syndrome or Turner Syndrome. BAV patients are of particular interest because this condition is not uncommon with a prevalence of about 1% in the general population [3–5]. However, BAV patients seem to be at relatively low risk for aortic dissection [6]. On the contrary, Marfan Syndrome has a lower prevalence of 6.5/100,000, but these patients are at high risk of acute aortic dissection [7].

The hemodynamic changes associated with exercise, and specifically the increase in blood pressure, is potentially associated with an enhanced risk of aortic growth and acute aortic dissection in the context of a thoracic aortic disease (TAD). Current guidelines state that patients with TAD should avoid strenuous resistance or isometric exercise and competitive sports [8–10]. Due to the lack of data, however, these European, Canadian and American guidelines are characterized by low levels of evidence [8–10]. Recommendations for specific

patient groups, such as patients with BAV, are in line with these guidelines. However, MFS patients are advised to only participate in low and moderate intensity sports with regular checks including echocardiography every 6 months, even if aortic root dilatation is absent [11–13].

The importance of daily exercise became clear in the 1950's when an inverse relationship between physical activity and cardiovascular risk was discovered [14]. Ever since, it has become well understood that a sedentary lifestyle is an important modifiable risk factor for cardiovascular disease and mortality [15]. Furthermore, regular exercise is known to prevent and reduce hypertension [16]. For TAD patients it is evenly important to not have a sedentary lifestyle, but also to prevent thoracic aortic growth and the occurrence of aortic dissection, creating a difficult paradox for clinicians. In this study, we sought to provide an up-to-date systematic review of the available evidence on exercise and sports participation in TAD patients including those with inherited aortopathies, and identify gaps in knowledge. We particularly aimed to find evidence on: (1) the aortic remodelling associated with regular exercise training and upper limits of dimensions in physically active individuals, (2) the risk of acute thoracic aortic dissection during exercise, and (3) the impact of exercise on the thoracic aorta in specific patient groups, especially in BAV and MFS patients.

Article highlights

- Strenuous and isometric exercise might contain a risk of causing acute aortic dissections in patients with TAD. However, the exact relationship between exercise and acute aortic dissections remains unclear.
- A potential positive effect of mild to moderate dynamic exercise training on thoracic aortic dilatation rate and aortic wall structure was reported in mice with MFS.
- There is a gap in knowledge about the positive and negative effects of exercise and sports participation in TAD patients. More research is needed to investigate the exact effects of different types of exercise on thoracic aortic dilatation growth rate and the risk of aortic dissection.

2. Methods**2.1. Literature search**

A broad systematic search was performed in Medline, Embase and Web of Science on 2 August 2018. The following search terms (including synonyms) were used: exercise, sports, athletes, training and thoracic aortic aneurysm, thoracic aortic dissection. Additionally, search terms were included for various inheritable connective tissue disorders: Marfan syndrome, Loeys-Dietz syndrome (including aneurysm osteoarthritis syndrome e.g. SMAD3 mutation), Turner syndrome, Ehlers-Danlos syndrome and bicuspid aortic valve. The exact search details are shown in supplemental file 1. Additional publications were obtained by hand searching, and reference lists were cross-checked to identify possible relevant papers overlooked by the original search. Duplicates were identified and removed.

2.2. Study selection

Titles and abstracts were screened for eligibility by two independent researchers (CT and LB). Only articles in the English language were included. Solely original data was included, therefore reviews and meta-analysis were excluded. Furthermore, book chapters, double publications on the same cohort and conference abstracts were excluded. Papers that could not be accessed in full text were also excluded. Only case reports on acute thoracic aortic dissection associated with exercise were included, while case reports on thoracic aortic dilatation and case reports on aortic dissections not related to exercise were excluded. Papers on thoracic aortic dilatation in athletes were only included if aortic diameters were reported. Of all potentially eligible papers the full text was reviewed. In case of disagreement a third reviewer was asked for counsel (JR) and eligibility was assessed by reasoning.

3. Results**3.1. Search results**

Figure 1 shows the flowchart of the study selection. Our search identified a total of 1652 unique publications. After reviewing the titles and abstracts 1530 papers were excluded, and 122 potentially eligible papers were reviewed in full text. Finally, 26 studies and 12 case reports were included. We

grouped the selected papers based on the abovementioned subjects of interest. Sixteen studies were found on thoracic aortic diameters in athletes. Three studies and twelve case reports were identified on the occurrence of acute aortic dissections during exercise. Three studies reported on exercise in MFS and four evaluated exercise in patients with BAV. Unfortunately, no papers were identified addressing the association between exercise and thoracic aortic dilatation or risk of dissection in patients with Loeys-Dietz syndrome, aneurysm-osteoarthritis Syndrome (AOS e.g. SMAD3 mutation), vascular Ehlers-Danlos syndrome or Turner syndrome.

3.2. Thoracic aortic dimensions in athletes

We identified 16 papers published between 1981 and 2015 which evaluate aortic dimensions in athletes practicing a variety of sports disciplines, shown in Table 1. Almost all papers were cross-sectional cohort studies (15/16), and one was a longitudinal cohort study. The number of included patients differed greatly, ranging from 9 to 1929 participants. Eleven studies compared aortic diameters in athletes to a sedentary control group, shown in Figure 2 [17–27]. Overall, outcomes of these studies show that athletes have significantly larger absolute aortic diameters than controls. However, the reported differences in absolute mean aortic root diameters are small: varying between 0.6 and 4 mm. Aortic diameter measurement was performed at the level of the aortic root in all studies, only two studies measured aortic diameter at multiple levels [23,28]. One study reported aortic root diameters corrected for body surface area (BSA) and found no significant difference between athletes and controls, although the absolute aortic root diameters were significantly different between the groups [26]. Three papers only included female athletes [17,19,20], and five papers only included male athletes [18,21–24]. The 99th percentile of aortic root dimensions in male athletes was found to be 40 mm and 34 mm for female athletes [29]. Five articles reported the prevalence of aortic root dilatation [27,29–32]. In these articles, different definitions of aortic dilatation were used, as shown in Table 1. The reported prevalence of aortic dilatation among athletes was low (0.26–1.3%), except in one cohort of athletes from the US national volleyball team, in which 6% of female athletes had an aortic root diameter ≥ 34 mm, and 8% of male athletes had an aortic root diameter ≥ 40 mm [32]. However, these volleyball players were very tall with an average body height of 198.2 ± 8.0 cm in males and 184.1 ± 7.4 cm in females.

Four studies evaluated differences in aortic diameter between strength trained and endurance trained athletes [21,27,28,31]. Three studies report a small but significant difference in absolute aortic root diameters, with slightly larger aortic root diameters in strength trained athletes than athletes who perform dynamic exercise. The mean differences reported ranged from 2.1–5 mm. However, mean aortic root measurements were all below 40 mm [27,28,31].

3.3. Exercise-related acute thoracic aortic dissections

Table 2 presents all case reports and case series reporting the occurrence of thoracic aortic dissections during exercise. The papers were published between 1987–2016 and each describe




Figure 1. Flowchart of literature search and selection of studies.

1 to 31 cases of acute thoracic aortic dissection occurring during sports activities. A total of 49 patients were described, of whom 42 suffered Stanford type A thoracic aortic dissections and 7 patients had Stanford type B dissections. Remarkably, only 2 out of 49 patients (4%) were female. The age ranged from 12 to 76 years. However, many reports only included young patients, with half of the papers reporting on patients up to 20 years of age [33–38]. In the majority of cases (26/49) weightlifting was the type of sport associated with the occurrence of aortic dissection [34–37,39–41]. MFS was diagnosed after presentation in four patients and one patient was known to have a connective tissue disorder other than MFS, which was not specified. Notably, family history was not obtained or reported in 7 of the 12 papers [34,35,39,41–44].

Furthermore, three retrospective cohort studies on sports related acute aortic dissections type A (AD-A) were identified, shown in Table 3. Only one paper focussed on the different types of sports practised during AD-A [45]. This study described 650 patients with a mean age of 62.3 years in patients with sports-associated AD-A and 63.7 years in non-sports associated AD-A. Of all AD-A's 4.1% was found to be associated with sports activities [45]. The type of sport most often reported was golf (32%), followed by swimming and cycling (each 16%), weight lifting (12%), and dancing (8%). Figure 3 illustrates the distribution of sports-

related AD-A's reported in this study over the different sports categories. These exercise related AD-A's occurred in all age groups and there was no significant difference in sex distribution between sports related AD-A's (60% males) and non-sports related AD-A's (52% males) [45]. Two retrospective cohort studies were identified specifically studying the occurrence of AD-A during a specific exercise: sexual intercourse and alpine skiing [46,47]. The first reports exercise and sexual intercourse associated AD-A's in a cohort of 365 patients and found a much higher percentage of 68% exercise associated AD-A's, with no significant difference between males and females. In this study, evidence of MFS was present in only 0.9% of patients. AD-A associated with sexual intercourse occurred only in males (17/245) [46]. The other retrospective cohort study by Schachner et al [47], reported on AD-A's occurring during winter season and they found that 22% of all AD-A's were associated with alpine skiing, and the majority of these cases were unrelated to trauma (82%). There was no significant difference in sex distribution between skiing associated AD-A (88% males) and AD-A not associated with skiing (77% males).

3.4. Exercise in patients with BAV

We identified four papers reporting on the association between exercise and thoracic aortic diameters or thoracic aortic

Table 1. Aortic dimensions in trained individuals.

Ref. nr.	First author	Year	Journal	Study design	n (total)	Study population			Outcome			
						Patient population	Sex (% female)	Control group	Outcome measures	Prevalence aortic dilatation	Results	Conclusion
17	Rubal	1981	Medicine and Science in Sports and Exercise	Cross-sectional cohort study	9	Female softball athletes	100	10 age- and body size-matched sedentary controls	Left ventricular parameters, aortic root diameter	Not reported	Mean aortic root diameter in athletes was 22 mm \pm 1 and 23 mm \pm 1 in controls.	There was no statistically significant difference in aortic root diameter between athletes and controls.
18	Wieling	1981	British Heart Journal	Longitudinal cohort study	23	Male oarsmen: seniors (n = 14) and freshmen (n = 9)	0	17 healthy age-matched controls	Left ventricular parameters, right ventricular diameter, aortic root diameter at end-diastole	Not reported	At the beginning of the season aortic root diameter was 31,3mm \pm 1,5 in senior oarsmen, 28,8 mm \pm 2,0 in freshmen, and 29,9 mm \pm 2,8 in control subjects.	At the beginning of the season aortic root diameter was greater in senior oarsmen than in freshmen, but did not differ from control subjects. No consistent increase was observed in aortic root diameter at the end of the season.
19	Crouse	1992	Research Quarterly for Exercise and Sport	Cross-sectional cohort study	15	Female basketball athletes	100	22 age-matched non-athletic controls	Left ventricular parameters and aortic root diameter	Not reported	Aortic root diameter was 25,7 mm \pm 3,3 in athletes and 21,7 mm \pm 2,4 in controls.	Aortic root diameter was significantly higher in athletes than in controls.
20	Pelliccia	1996	JAMA	Cross-sectional cohort study	600	Female athletes from Italian national teams	100	65 age-matched untrained females and 738 age-, ethnicity, sports discipline and training intensity matched elite male athletes	Left ventricular parameters, aortic root diameter	Not reported	The aortic root was 27,6 mm \pm 2,5 in athletes, 27,0 mm \pm 1,8 in controls, and 30,3 mm \pm 2,0 in male athletes.	The aortic root was significantly larger (2% larger) in athletes than in controls, and significantly larger in male athletes.
21	Whyte	1999	International Journal of Sports Medicine	Cross-sectional cohort study	29	Elite modern pentathletes (n = 11) and triathletes (n = 18)	0	13 sedentary controls	Left ventricular parameters, aortic annulus diameter	Not reported	Aortic annulus diameter was 28,8 mm \pm 5,1 in triathletes, 33,5 mm \pm 2,1 in pentathletes and 31,1 mm \pm 1,9 in controls.	There was no statistically significant difference in aortic annulus diameter between triathletes, pentathletes and controls. Whilst the large training volume elicits significant morphological adaptation in triathletes and modern pentathletes, all measures were within normal limits.
30	Kinoshita	2000	American Heart Journal	Cross-sectional cohort study	1929	Athletes active in competitive sports	19	No control group	Aortic root diameter	Aortic root >40 mm: 0,26–0,36% in basketball and volleyball players	0,26–0,36% of athletes had aortic root dilatation >40 mm, 0,96% of basketball en volleyball players had aortic root dilatation >40 mm.	A higher prevalence of aortic dilatation is to be anticipated among basketball and volleyball players, many of whom are very tall.
22	Dzudie	2006	European Journal of Echocardiography	Cross-sectional cohort study	21	Male handball players	0	21 age-, sex-, height- and weight-matched sedentary men	Left ventricular parameters and aortic root diameter	Not reported	Aortic root diameter was 29,6 mm \pm 3,6 in control subjects and 30,3 mm \pm 2,8 in handball players.	Aortic root diameter was slightly larger in control subjects than in handball players, but this difference was not significant.

(Continued)

Table 1. (Continued).

Ref. nr.	First author	Year	Journal	Study design	Study population			Outcome				
					n (total)	Patient population	Sex (% female)	Control group	Outcome measures	Prevalence aortic dilatation	Results	Conclusion
23	Babaee Bigi	2007	American Journal of Cardiology	Cross-sectional cohort study	100	Male elite athletes	0	128 age- and height-matched healthy men	Aortic diameter, aortic regurgitation	Not reported	Aortic diameters were measured at Ann 25.1 ± 2.9 in athletes and 21.8 ± 2.4 in controls, Sov 38.2 ± 4.1 in athletes and 31.6 ± 3.2 in controls, STJ 34.1 ± 2.8 and 29.5 ± 3.1 in controls, AA 36.1 ± 4.5 in athletes and 31.0 ± 2.9 in controls.	Aortic root diameters in all segments of the aortic root were significantly greater in elite strength-trained athletes compared with an age- and height-matched population.
29	Pelliccia	2010	Circulation	Cross-sectional cohort study	2317	Highly trained athletes	44	No control group	Aortic diameter	Aortic root >40 mm: 1.3% of male athletes. Aortic root >34 mm: 0.9% of female athletes	Mean aortic root diameter was 32.2 mm ± 2.7 in male athletes with 99th percentile 40 mm, and 27.6 mm ± 2.6 in female athletes with 99th percentile 34 mm.	Aortic root enlargement (40 mm in males and 34 mm in females) is particularly uncommon in highly trained athletes.
31	D'Andrea	2010	American Journal of Cardiology	Cross-sectional cohort study	615	Elite athletes: endurance-trained athletes (n = 370) and strength-trained athletes (n = 245)	33	No control group	Aortic diameter	Ascending aortic dilatation >95% CI of overall distribution: 1% of athletes	The mean aortic root diameter at the Sov was 31 mm (2.8–3.6) in endurance trained athletes and 36 mm (3.2–4.2) in strength trained athletes.	The aortic root diameters at all levels were significantly greater in strength trained athletes. Significant ascending aortic dilatation and aortic regurgitation proved to be uncommon in strength trained athletes.
24	Carlsson	2010	European Journal of applied physiology	Cross-sectional cohort study	10	Male endurance athletes	0	10 untrained men	Cardiac functional parameters, cardiac structural parameters (among which aortic root diameter)	Not reported	Aortic root diameter was 30.4 mm ± 3.2 in athletes and 28.4 mm ± 4.4 in controls.	Aortic root diameter was not significantly larger in athletes than in controls.
25	Caselli	2011	European Journal of Echocardiography	Cross-sectional cohort study	429	Athletes from Italian national teams	22	98 healthy controls	Left ventricular parameters and aortic root diameter	Not reported	Aortic root diameter was 31.0 mm ± 3.8 in athletes and 28.7 mm ± 3.3 in controls.	Aortic root diameter was significantly larger in athletes than in controls.
26	Krol	2011	Echocardiography	Cross-sectional cohort study	38	Members of the Polish Olympic team (rowing, cycling, speed-skating)	16	41 sex and age matched, healthy sedentary individuals	Left ventricular parameters, right ventricular parameters, aortic diameter (level of measurement not specified)	Not reported	Aortic diameters were 33 mm ± 4 in het athletes and 29 mm ± 2 in controls. After indexing for BSA the mean aortic size index for athletes was 1.6 cm/m ² ± 0.1 and 1.5 cm/m ² ± 0.2 for controls.	Aortic diameters were significantly larger in het athletes group than in controls. However after indexing for BSA there was no significant difference between the groups.

(Continued)

Table 1. (Continued).

Ref. nr.	First author	Year	Journal	Study design	n (total)	Study population			Outcome		
						Sex (% female)	Control group	Outcome measures	Prevalence aortic dilatation	Results	Conclusion
27	D'Andrea	2012	Journal of the American Society of Echocardiography	Cross-sectional cohort study	410	38	240 healthy controls	Aortic root diameter, aortic root distensibility and elasticity	Aortic root dilatation >95% CI of overall distribution: 1% of male power athletes	Aortic root diameter was 36 mm ± 5 in strength trained, 31 mm ± 6 in endurance trained athletes and 32 mm ± 3 in controls.	Aortic root diameters and stiffness were significantly greater in strength trained athletes than in endurance trained athletes and controls. While aortic distensibility was higher in endurance trained athletes compared to controls.
28	Aparci	2013	Experimental & Clinical Cardiology	Cross-sectional cohort study	60	Unknown	No control group	Aortic diameters, left ventricular parameters, left atrial diameters	Subjects with abnormally enlarged aortic diameters (≥40 mm) were excluded	In the strenuous activity training group mean aortic diameters were: 35.6 mm (SoV) and 36.8 mm (AA), in the ordinary activity training group the mean aortic diameter was 33.5 mm (SoV) and 34.4 mm (AA).	Aortic root and ascending aortic diameter were significantly higher in the strenuous activity training group.
32	Davis	2015	Clinical Journal of Sport Medicine	Cross-sectional cohort study	70	47	No control group	Aortic diameter and Ghent criteria (signs of Marfan syndrome)	Aortic root diameter ≥40 mm: 8% of male athletes. Aortic root diameter ≥34 mm: 6% of female athletes	34% of the athletes had at least 1 characteristic of MFS (Ghent criteria) but none had more than 2 characteristics.	Elite US volleyball players have a higher than expected prevalence of dilation of the aortic sinuses and ascending aorta. In the absence of MFS.

Ann; Aortic Annulus, SoV; Sinus of Valsalva, STJ; Sinotubular Junction, AA; Ascending Aorta




Figure 2. Studies that reported aortic root diameters in athletes and controls.

* This study reported only aortic annulus diameter, not aortic root diameter

dilatation rate in BAV patients, which are shown in Table 4. Of these, three papers came from the same research group [48–50]. Two of which compared athletes with BAV to athletes with a normal tricuspid aortic valve (TAV) [48,49]. Both reported significantly larger aortic diameters at all measured levels in BAV athletes compared to TAV athletes. However, all reported mean diameters were below 36 mm. One cross-sectional study, which included 58 competitive athletes with BAV, showed no correlation between aortic dimensions and duration of training [48]. Two longitudinal studies presented by the same research group reported on mean aortic diameter growth rate in BAV athletes, presumably describing the same patients. The mean growth rates reported were: 0.78 mm/year at the aortic annulus (Ann), 0.61 mm/year at the Sinuses of Valsalva (SoV), 0.81 mm/year at the sinotubular junction (STJ) and 0.98 mm/year at the proximal ascending aorta (AA) [49,50]. The mean age of these two cohorts of BAV athletes were 19 ± 8.8 years and 25 ± 11 years. No significant increase of aortic diameter was reported in TAV athletes (mean age 25 ± 5 years) after five years of follow-up. Another longitudinal study by Spataro et al. found no clear association between sports participation and valve deterioration in BAV patients, with a mean follow-up duration 13 years [51]. Unfortunately, no aortic diameter measurements were reported and no conclusions can be drawn about the effect of exercise on the aortic diameter in this cohort of BAV athletes. Only one paper compared BAV athletes to sedentary BAV subjects. This article reported no difference in aortic growth rate between the two groups at all measured levels of the thoracic aorta: Ann, SoV, STJ and AA [50].

3.5. Exercise in marfan syndrome

Table 4 presents the three papers on exercise in MFS, all published in 2017. Two papers describe mouse model studies investigating the effects of mild-moderate dynamic exercise on the aortic wall in MFS mice [52,53]. Both were controlled trials with one or more dynamic exercise training groups and a sedentary group. The follow-up duration of both studies was five months. Both papers reported a reduction of aortic diameter growth rate in MFS mice performing mild to moderate dynamic exercise compared to sedentary MFS mice [52,53]. Also, in mice performing dynamic exercise, the aortic wall became stronger compared to sedentary

MFS mice. This was testified by the larger amount of mechanical stress on the aorta required to induce rupture of the aortic wall [52]. Exercise seemed to improve aortic wall elasticity in one study [52], but no significant improvement was found in the other [53]. Dynamic exercise was not found to increase lamina ruptures, indicating no additional structural damage in the tunica media [53]. An optimum of protective effects was found at a training intensity level of 55–65% of maximum oxygen uptake (VO₂max), while higher intensity of dynamic exercise training seems to blunt the positive effects [52]. The third paper is a small prospective cohort study that evaluated the feasibility and effects of a three-week rehabilitation program in 19 MFS patients with a mean age of 46.7 ± 7.8 years [54]. During the one-year follow-up, no adverse medical events were reported, physical fitness improved, and psychological distress decreased. These effects were already present after three weeks of rehabilitation, and mostly remained persistent throughout the one-year follow-up [54]. Unfortunately, no information on aortic diameters was provided.

4. Discussion

To our knowledge, this is the first systematic review describing the effect of exercise and sports participation in TAD patients. We were not able to identify any controlled or randomized trials evaluating the longitudinal effect of exercise on survival or risk of aortic dissection in TAD patients. When focusing on the association between exercise and thoracic aortic growth rate very limited data can be found. In total, we identified 38 papers of interest, of which 9 were case reports, 3 case series, 8 longitudinal cohort studies and 16 cross-sectional cohort studies. Two were mouse-model studies: one non-randomized controlled trial and one randomized controlled trial. Assessment of methodological quality of the included papers was planned, but ultimately not performed quantitatively, since the large variety of study designs made consistent and comparable quality assessment impossible and meaningless. Eventually it can be concluded that most papers would reach low scores.

4.1. Are aortic dimensions different in athletes?

Screening before participating in competitive sports provides a lot of easily accessible data resulting in a large number of

Table 2. Case reports and case series on acute aortic dissections during exercise.

Ref. nr.	First author	Year	Journal	Study design	Study population			Outcome		Aortopathy	Family history	Conclusion
					n (total)	Age (years)	Sex	Stanford Classification	Max. aortic diameter			
33	Bain	1987	The American Journal of Forensic Medicine and Pathology	Case-report	1	20	M	Type A	60 mm	Fitness	Positive for sudden death	This case demonstrates Marfan Syndrome presenting as sudden, unexpected death.
39	De Virgilio	1990	The Annals of Thoracic Surgery	Case series	4	22–57	M	Type A	Unknown	Weightlifting	Unknown	Individuals who have evidence of cystic medial disease or family histories of this disease should avoid weight lifting.
34	Schor	1993	Journal of Vascular Surgery	Case-report	1	18	M	Type B (periaortic hematoma)	Unknown	Weightlifting	Unknown	Weight lifting, with its profound cardiovascular effects, may be the major, if not sole cause of aortic dissection.
35	Baumgartner	1997	The Annals of Thoracic Surgery	Case-report	1	19	M	Type A	Unknown	Weightlifting	Unknown	Individuals with Marfan Syndrome, cystic medial disease or family histories of the disorder should be strongly urged to refrain from weight-lifting activities.
36	Eleferiades	2003	Journal of the American Medical Association	Case series	5	19–53	Unknown	Type A	40–52 mm	Weightlifting, push ups, heavy lifting	Positive for aortic disease in 1 patient	The risk of weight lifting as a cause of aortic dissection has generally been underappreciated.
40	Hogan	2005	Emergency Medicine Journal	Case-report	1	27	M	Type A	Unknown	Weightlifting	Positive for aortic dissection	Aortic dissection should be considered in symptomatic patients with a family history of early cardiac deaths, suspect of a connective tissue disorder, or who practice weightlifting.
37	Hatzaras	2007	Cardiology	Case series	31	19–76	30 M, 1 F	Type A: 87% (n = 27), Type B: 13% (n = 4)	30–78 mm	Weightlifting or push-ups (n = 16), heavy lifting (n = 9), dynamic exercise like swimming or tennis or racquetball (n = 5)	Positive for aortic disease in 3 patients	Increased blood pressure due to heavy weight lifting raises aortic wall stress to a level that produces aortic dissection in individuals with pre-existing mild to moderate aortic enlargement.
38	Uchida	2009	Interactive CardioVascular and Thoracic Surgery	Case-report	1	12	M	Type B	Unknown	Swimming	Negative	Swimming coaches and pediatricians should recognize that swimming exercises like the butterfly stroke are a risk factor for aortic dissection in children.
42	Westaby	2011	Circulation	Case-report	1	28	M	Type B	Unknown	Soccer	Unknown	Aortopathy associated with a bicuspid aortic valve and coarctation may contribute to aneurysmal transformation and rupture.
43	Chattranukulchai	2013	British Medical Journal Case Reports	Case-report	1	38	M	Type A	52 mm	Bowling	Unknown	Early diagnosis of Marfan Syndrome is crucial as it has a positive influence on the outcome.
41	Ozyildirim	2015	The American Journal of Cardiology	Case-report	1	28	M	Type A	Unknown	Weightlifting	Unknown	Weight lifting creates significant stress along the aortic wall and this produces predisposition to acute aortic dissection.
44	Cereda	2016	European Heart Journal	Case-report	1	25	F	Type A	100 mm	Volleyball	Unknown	Echocardiography has a potential role in preventing tragic sudden death in sport.

Table 3. Papers on sports related aortic dissections.

Ref. nr.	First author	Year	Journal	Study design	Study population			Outcome			
					n (total)	Patient population	Sex (% female)	Definition	Missing (%)	Exercise related dissections (%)	Additional results
47	Schachner	2013	BioMed Research International	Retrospective cohort study	140, of which 77 (55%) during winter season	Patients with aortic dissection involving the ascending aorta during the winter season (from the beginning of November until the end of April).	21 (16/77)	Onset of symptoms during alpine skiing.	Not reported	22	In 14/17 (82%) patients symptoms occurred during recreational skiing without additional trauma. Only 1 patient (6%) had a skiing accident with consequent ascending aortic dissection.
46	Gansera	2015	The Thoracic and Cardiovascular Surgeon	Retrospective cohort study	365	Patients who underwent surgery for aortic dissection involving the ascending aorta.	32	Onset of symptoms occurred during physical exercises, such as sports or lifting of heavy weights.	24	68	Peri- or postcoital aortic dissection occurred in none of the females, but in 17 of 245 males. Mortality in females < 65 years was higher (20.2%), compared with their male counterparts (14.9%). The sport most frequently associated with type A aortic dissections in this series was golf (n = 8; 32%), followed by swimming and cycling (each n = 4; 16%), then weight lifting (n = 3; 12%), dance (n = 2; 8%), and finally, long-distance running, fencing, table tennis, and archery, (each n = 1; 4%).
45	Itagaki	2017	Surgery Today	Retrospective cohort study	650	Patients who underwent surgery for aortic dissection involving the ascending aorta.	47 (323/615)	Onset of symptoms during a sports activity. Non-sports exertion, such as lifting or moving a heavy load, defecation, or sexual activity, were classified into the non-sports group.	5	4.1 (5% of all daytime type A dissections)	

studies performed in athletes. The absolute thoracic aortic diameters reported are larger in athletes compared to non-athletic controls, although the differences are very small. This is consistent with the findings of a fairly recent meta-analysis reporting a small but significantly larger aortic root diameter in elite athletes than in non-athletic controls [55]. One study showed that after indexing aortic root diameter for BSA, the significant difference in aortic root diameter between athletes and controls was blunted [26]. Indicating the differences in aortic diameter might be attributed to a difference in body size between athletes and controls, which is known to be associated with thoracic aortic diameter [56]. The larger aortic diameters reported in athletes are therefore not necessarily caused by a pathological process, but presumably result from higher cardiac output and difference in body size. Whether this is associated with an increased risk of aortic dissection is yet to be determined. The findings mentioned above seem to be comparable in male and female athletes. However, they might not be applicable to less intensively trained and older individuals, and patients who already have TAD.

4.2. Is there an association between exercise and acute thoracic aortic dissections?

The incidence of sudden cardiac death among the younger population (< 40 years) is approximately 1.3 to 8.5 per 100,000 person-years [57]. Approximately 1–5% of sudden death in young athletes is caused by acute aortic dissections [58–60]. Over the past decades, there have been many reports, especially case-reports, that link exercise (mainly high intensity static exercise) to acute aortic dissection. Aortic dissection is an emergency situation with a reported in-hospital mortality of up to 33% [61]. This has made clinicians cautious when counselling TAD patients about exercise. We found a striking difference in the reported amount of acute thoracic aortic dissections related to exercise: Itagaki et al. [45] reported a relation to exercise in 4.1% of all AD-A's and Gansera et al. [46] in 68% of all AD-A's. This difference might partially be explained by a different definition of 'sports related aortic dissections'. Whereas one study classified non-sports exertion and Valsalva manoeuvres such as lifting or moving a heavy load, defecation, or sexual activity, into the non-sports group (Itagaki et al.), the other classified these into the sports group (Gansera et al.). The true amount of sports-related type A aortic dissections might be somewhere in between, such as the 22% of all winter season AD-A's related to alpine skiing, as reported by Schachner et al [47]. Apart from the theoretical physiological impact, we found no evidence supporting the theory of static exercise being more prone to inducing acute thoracic aortic dissections than dynamic exercise. The majority of the identified case reports described acute thoracic aortic dissection during weightlifting. However, this might be due to selection bias, since a larger series showed the type of sport associated with AD-A most frequently was golf (32%), which is classified as a low isometric and low isotonic sport (skill category, Figure 3) [45]. It seems that aortic dissection can also occur during relatively mild intensity sports such as golf, but for interpreting the results of Figure 3 keep in mind that this is based on only one study and more research is clearly warranted. Of course, the size and composition of the study



Figure 3. Classification of sports and sports related aortic dissections.

Most sports require a combination static and dynamic exercise. In order to facilitate counselling about sports participation this simplified classification of the most common olympic sports disciplines was created, according to the relative isometric and isotonic components of exercise and resulting cardiovascular adaptation. Underneath the distribution of sports related aortic dissections type A (AD-A) over the four categories is displayed, based on data published by Itagaki et al. [45]. The classification of sports was reprinted from: A. Pelliccia; S. Caselli, European Association of Preventive Cardiology (EAPC) and European Association of Cardiovascular Imaging (EACVI) joint position statement: recommendations for the indication and interpretation of cardiovascular imaging in the evaluation of the athlete's heart, European Heart Journal, 2017, Volume 39, Issue 21, Pages 1949–1969, by permission of Oxford University Press. AD-A = Aortic Dissection Stanford type A

population is crucial here. When a study includes all patients with dissection, the mean age will be relatively high, and golf will be a sport which is prevalently practised. In younger cohorts, a totally different sports involvement pattern is likely to be found. Furthermore, this study was conducted in Japan where golf is known to be a very popular form of exercise. Therefore, in the absence of reliable information on rates of sports participation, no conclusion can be drawn on the association between dissection and a specific sports activity. Concerning the impact of sex, it was striking that almost all case reports describe males with acute aortic dissections related to exercise. However, this does not seem representative since the larger series both from Itagaki and Gansera reported no significant differences in sex distribution [45,46]. Population based studies might provide additional information, but data on the prevalence of participation in different types of sports, aortic diameters and long-term follow-up are scarce.

4.3. How does exercise influence the thoracic aorta in BAV and marfan patients?

Although bicuspid aortic valve is the most prevalent congenital heart disease and an important underlying etiology of thoracic aortic dilatation, the association between exercise and aortic diameter and growth rate has been investigated to a limited extent in this patient group. In BAV patients,

Galanti et al. and Stefani et al. state that the aortic growth rate they reported in BAV athletes does not differ from aortic growth rate reported in the general BAV population [49,50]. Indeed, the reported dilatation rate of 0.98 mm/year found in the athletes with BAV seems comparable to the reported aortic dilatation rate in various studies reporting aortic growth rate in the general BAV population [62,63]. Even though the BAV populations studied by Stefani et al and Galanti et al were relatively young (19 ± 8.8 years and 25 ± 11 years), and younger age is known to be associated with higher aortic growth rates [64]. These findings suggest that aortic growth rate is not significantly influenced by exercise in BAV patients.

Two recently published papers have investigated the effect of dynamic exercise on the thoracic aorta in mice with MFS. Both studies reported mild to moderate dynamic exercise had a positive effect on aortic growth rate and seemed to improve aortic wall structure. This suggests that exercise does not only have potential negative effects on the thoracic aorta in TAD patients, but might be actually be beneficial [52,53]. Further research is needed to evaluate these potential positive effects of exercise on the thoracic aorta in MFS patients and patients with other thoracic aortic diseases. Especially, since mouse model studies on the aorta might not always be reliable [65]. One randomized trial has been performed in patients with an abdominal aortic aneurysm (AAA) in 2014 [66]. In this trial 140 patients with AAA

Table 4. Papers on exercise in patients with Marfan Syndrome and bicuspid aortic valve.

Ref. nr.	Group	First author	Year	Journal	Study population			Outcome						
					Study design	n (total)	Patient population	Genetic mutation	Sex (% Female)	Control group	Follow-up	Results	Conclusion	
51	BAV	Spataro	2008	International Journal of Sports Medicine	Longitudinal cohort study	81	Competitive athletes with BAV from Italian national teams divided into 2 groups: the low-risk group (n = 51) and the high-risk group (n = 30)	NA	10	No control group	13 ± 4.9 years (range 5–19 years)	Aortic regurgitation, aortic stenosis and left ventricular parameters, aortic root diameters.	Over the follow-up period, six of the initially low-risk athletes (7%) and all of the high-risk patients showed significant worsening of morphologic features of bicuspid aortic valve and/or incidence of symptoms. In high risk subjects the progression of valvular disease occurred independently from the former athletic activity.	Continued sport participation is not responsible itself of BAV worsening. However, long-term athletic training may be associated with progressive worsening of the valvular lesion and the appearance of clinical symptoms.
48	BAV	Stefani	2008	British Journal of Sports Medicine	Cross-sectional cohort study	58	Non-elite but competitive athletes with BAV	NA	0	75 non-elite but competitive athletes with TAV	NA	Aortic regurgitation, aortic stenosis and aortic root diameters.	Aortic root dimensions at all levels were significantly greater in athletes with BAV than in athletes with a normal TAV. No relation was found with age, body surface area, aortic regurgitation or years of training.	A short echocardiographic examination should be performed at least once during an athlete's sporting life.
49	BAV	Galanti	2010	British Journal of Sports Medicine	Longitudinal cohort study	88	Athletes with BAV and mild aortic regurgitation	NA	Unknown	56 athletes with TAV	5 years (30/88 subjects)	Left ventricle parameters and aortic diameters	There was a progressive increase at each measured aortic level (Ann: 0.78 mm/year; Sov: 0.61 mm/year; ST: 0.81 mm/year; AA: 0.98 mm/year). This increase became significant from the last 2 years of the 5-year follow up. In TAV athletes, there was no significant increase of aortic diameters (Ann 0.17 mm/year; Sov 0.12 mm/year; ST 0.21 mm/year; AA: 0.32 mm/year)	In athletes with BAV, aortic dimensions increase significantly more than in TAV athletes, but do not differ from those in the general BAV population.
50	BAV	Stefani	2014	Cardiology Research and Practice	Longitudinal cohort study	292	Subjects with BAV who were evaluated at the Sports Medicine and Exercise Centre, divided into three different groups: athletes (n = 210), sedentary (n = 59), and ex-athletes (n = 23)	NA	Unknown	No control group	5 years	BAV morphology classification, left ventricular parameters and aortic diameters	Typical BAV morphology was most frequent in all three groups (68% athletes, 67% sedentaries, and 63% ex-athletes). The aortic dimensions showed a progressive enlargement during follow-up, with no difference between athletes and sedentary subjects. There was a progressive increase at each measured aortic level (Ann: 0.78 mm/year; Sov: 0.61 mm/year; ST: 0.81 mm/year; AA: 0.98 mm/year).	In BAV patients the ascending aorta is involved in normal progressive, not necessarily pathological, enlargement.

(Continued)

Table 4. (Continued).

Ref. nr.	Group	First author	Year	Journal	Study population			Genetic mutation	Sex (% female)	Control group	Outcome			Conclusion
					Study design	n (total)	Patient population				Outcome measures	Follow-up	Results	
54	MFS	Benninghoven	2017	Orphanet Journal of Rare Diseases	Longitudinal cohort study	19	Patients with MFS or similar syndrome in stable condition	Not specified	71	No control group	Adverse events, physical fitness and psychological assessment	1 year	No adverse medical events were reported. Physical fitness improved from admission to discharge. Psychological distress decreased. Admission to discharge effects mostly persisted through the one year follow-up but declined to smaller sizes.	The three week rehabilitation program improved physical fitness and psychological wellbeing. Medical assessments ruled out adverse events caused by participation in the program.
52	MFS	Gibson*	2017	Journal of Applied Physiology	Non-randomised controlled trial (mouse-model study)	16	Male mice with MFS divided in a sedentary group (n = 10), voluntary exercise group (n = 3) and forced exercise group (n = 3)	Fbn1C1039G/+	0	19 mice without MFS, divided in a sedentary group (n = 4), voluntary exercise group (n = 7) and forced exercise group (n = 8)	Historical characteristics, isometric force, aortic wall elasticity and stiffness and combined benefit score for: elastin fiber length, elastin fragmentation, and elasticity	5 months	Both voluntary and forced exercise routines reduced aortic diameter, prevented aortic wall weakening, increased the breaking stress and improved aortic wall elasticity in MFS aorta. There is an optimum of protective effects at training intensity levels between 55% and 65% (of VO2max) which significantly reduces elastin fragmentation and disorganization within the aortic wall.	The present study provides helpful insights into the potential protective effects of a mild exercise routine in MFS patients in the absence of pharmacological interventions.
53	MFS	Mas-Stachurska*	2017	Journal of the American Heart Association	Randomized controlled trial (mouse-model study)	19	Mice with MFS, randomized to a sedentary group n = 9 and exercise group n = 10	Fbn1C1039G/+	47	21 mice without MFS, randomized to a sedentary group n = 11 and exercise group n = 10	Left ventricular parameters, aortic root diameter, aortic pulsatility, aortic stiffness and histological characteristics of aortic wall tissue	5 months	In MFS mice subjected to exercise aortic root diameter was smaller than in their sedentary littermates and aortic dilatation rate was blunted, becoming comparable to the controls. Exercise training improved aortic stiffness in control but not in MFS mice, but did not increase lamina ruptures in MFS mice, indicating no additional structural damage in the tunica media.	Moderate dynamic exercise prevented aortic root dilation and mitigates cardiac hypertrophy.

* Mouse-model study

MFS; Marfan Syndrome, BAV; Bicuspid Aortic Valve, TAV; Tricuspid Aortic Valve, Ann; Aortic Annulus, Sov; Sinus of Valsalva, STJ; Sinotubular Junction, AA; Ascending Aorta

were randomized to either standard care or exercise training including dynamic as well as isometric exercise (rowing). No difference in abdominal aortic growth rate was reported between the groups. Although AAA has a different aetiology than thoracic aortic aneurysms and should therefore be seen as a different disease entity, the findings of this study are promising. A randomized study, such as the one illustrated above for AAA patients, would provide important additional information about the effect of exercise in TAD patients.

5. Conclusion

Although several case reports have described aortic dissection occurring during exercise, no high-quality studies have been performed to illuminate the association between exercise and acute aortic dissection. In athletes, aortic diameters are only slightly larger than in controls. Evenly, aortic diameter growth rate does not seem to be enhanced by exercise in BAV patients. In mice with MFS a positive effect of mild to moderate dynamic exercise on the thoracic aorta diameter was




Figure 4. Hemodynamic response to exercise. **(a)** Response to dynamic exercise of progressively increasing workload. This causes a volume overload as a result of increased cardiac output (Q) and arterial blood pressure (ABP), with a decrease in total peripheral resistance (TPR). **(b)** Response to a static handgrip contraction. This causes a pressure overload as a result of increased blood pressure, but no decrease in total peripheral resistance. ABP (mm Hg): systolic, mean and diastolic arterial blood pressures; HR: heart rate (beats/min); Q: cardiac output (liters/min); SV: stroke volume (ml/beat); TPR: total peripheral resistance (PRU); VO2: oxygen uptake (ml/min/kg). Reprinted by permission from J.H. Mitchell and P.B. Raven, 'Cardiovascular Adaptation to Physical Activity,' in *Physical Activity, Fitness, and Health: International Proceedings and Consensus Statement*, edited by C. Bouchard, R.J. Shephard, and T. Stephens (Champaign, IL: Human Kinetics, 1994), 288.

found. There clearly is a gap in knowledge about the effects of exercise and sports participation in TAD patients. Currently there is no unequivocal evidence to support discouragement of exercise and sports participation in TAD patients. Hence, mild to moderate regular exercise should be encouraged, for its known positive effects on overall health. However, based on theoretical knowledge, participation in heavy static exercise should likely be avoided in TAD patients.

6. Expert opinion

When a patient is diagnosed with TAD, discussing lifestyle modification is mandatory. Next to cessation of smoking, controlling hypertension and prevention of obesity, discussing exercise and sports participation is important. However, there is not enough evidence to strongly discourage exercise or recommend any particular type of exercise or sport. Theoretically, high blood pressure is unfavourable. Therefore, it is important to distinguish between dynamic (also isotonic) and static (also isometric) exercise [61], since both initiate a different hemodynamic response (illustrated in Figure 4). On the other hand exercise and sports participation are also known to have many positive effects on cardiovascular and overall health. For the general population the Dutch and American health councils, as well as the World Health Organization recommend a target rate of 150 to 300 minutes per week of moderate to heavy intensity exercise [67–69], as participation in regular physical activity has shown to have many benefits [67,68].

Therefore, we believe it is mandatory to explain both the negative and positive effects of exercise to TAD patients. In order to create full understanding and ideally reach a shared decision, rather than imposing restrictions on sports participation. In order to prevent TAD patients becoming scared of physical activities and to minimize concerns, stress and anxiety further affecting TAD patient's quality of life, which was shown to be reduced compared to healthy controls [70].

Ideally future research would be (randomized) controlled trials longitudinally evaluating the effect of exercise on thoracic aortic aneurysm dilatation rate, the risk of thoracic aortic dissections, quality of life and survival of TAD patients. Secondly, the effect of different types and intensities of exercise on thoracic aortic growth rate acceleration needs to be evaluated. More research is especially needed in patients with Loeys-Dietz syndrome, aneurysm-osteoarthritis Syndrome (AOS e.g. SMAD3 mutation), vascular Ehlers-Danlos syndrome and Turner syndrome, on which we found no evidence at all.

7. Five-year view

In the upcoming five years we envision that more research will be carried out on the association between exercise and thoracic aortic growth and acute aortic dissection. Further exploring the potential beneficial effect of dynamic exercise on the aortic wall in humans is warranted. This knowledge will enable us to better understand and predict the risks of exercise and sports participation in TAD patients. This will hopefully enforce better counselling, with more detailed and well-founded advice to TAD patients.

Acknowledgments

The authors would like to acknowledge Gerdien B de Jonge.

Funding

This paper was funded by ZonMW (project number: 849200014) and the Dutch Heart Foundation (contract grant number: 2013T093).

Declaration of interest

The authors have no relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript. This includes employment, consultancies, honoraria, stock ownership or options, expert testimony, grants or patents received or pending, or royalties.

Reviewer disclosures

Peer reviewers on this manuscript have no relevant financial or other relationships to disclose.

ORCID

Carlijn G. E. Thijssen  <http://orcid.org/0000-0002-8101-5954>

Jolien W. Roos-Hesselink  <http://orcid.org/0000-0002-6770-3830>

References

Papers of special note have been highlighted as either of interest (*) or of considerable interest (***) to readers.

1. Olsson C, Thelin S, Stahle E, et al. Thoracic aortic aneurysm and dissection: increasing prevalence and improved outcomes reported in a nationwide population-based study of more than 14,000 cases from 1987 to 2002. *Circulation*. 2006;114(24):2611–2618.
2. Albornoz G, Coady MA, Roberts M, et al. Familial thoracic aortic aneurysms and dissections—incidence, modes of inheritance, and phenotypic patterns. *Ann Thorac Surg*. 2006;82(4):1400–1405.
3. Nistri S, Basso C, Marzari C, et al. Frequency of bicuspid aortic valve in young male conscripts by echocardiogram. *Am J Cardiol*. 2005;96(5):718–721.
4. Tutar E, Ekici F, Atalay S, et al. The prevalence of bicuspid aortic valve in newborns by echocardiographic screening. *Am Heart J*. 2005;150(3):513–515.
5. Movahed MR, Hepner AD, Ahmadi-Kashani M. Echocardiographic prevalence of bicuspid aortic valve in the population. *Heart Lung Circul*. 2006;15(5):297–299.
6. Girdauskas E, Disha K, Borger MA, et al. Risk of proximal aortic dissection in patients with bicuspid aortic valve: how to address this controversy? *Interact Cardiovasc Thorac Surg*. 2014;18(3):355–359.
7. Groth KA, Hove H, Kyhl K, et al. Prevalence, incidence, and age at diagnosis in Marfan Syndrome. *Orphanet J Rare Dis*. 2015;10:153.
8. Boodhwani M, Andelfinger G, Leipsic J, et al. Canadian cardiovascular society position statement on the management of thoracic aortic disease. *Can J Cardiol*. 2014;30(6):577–589.
9. Hiratzka LF, Bakris GL, Beckman JA, et al. 2010 ACCF/AHA/AATS/ACR/ASA/SCA/SCAI/SIR/STS/SVM guidelines for the diagnosis and management of patients with thoracic aortic disease. A report of the American college of cardiology foundation/American heart association task force on practice guidelines, American association for thoracic surgery, American college of radiology, American stroke association, society of cardiovascular anesthesiologists, society for cardiovascular angiography and interventions, society of interventional radiology, society of thoracic surgeons, and society for vascular medicine. *J Am Coll Cardiol*. 2010;55(14):e27–e129.
10. Erbel R, Aboyans V, Boileau C, et al. 2014 ESC Guidelines on the diagnosis and treatment of aortic diseases: document covering

- acute and chronic aortic diseases of the thoracic and abdominal aorta of the adult. The task force for the diagnosis and treatment of aortic diseases of the European Society of Cardiology (ESC). *Eur Heart J*. 2014;35(41):2873–2926.
11. Firoozi S, Sharma S, McKenna WJ. Risk of competitive sport in young athletes with heart disease. *Heart*. 2003;89(7):710–714.
 12. Pelliccia A, Zipes DP, Maron BJ. Bethesda conference #36 and the European society of cardiology consensus recommendations revisited. A comparison of U.S. and European criteria for eligibility and disqualification of competitive athletes with cardiovascular abnormalities. *J Am Coll Cardiol*. 2008;52(24):1990–1996.
 13. Braverman AC, Harris KM, Kovacs RJ, et al. Eligibility and disqualification recommendations for competitive athletes with cardiovascular abnormalities: task force 7: aortic diseases, including Marfan syndrome: a scientific statement from the American heart association and American college of cardiology. *J Am Coll Cardiol*. 2015;66(21):2398–2405.
 14. Morris JN, Heady JA, Raffle PA, et al. Coronary heart-disease and physical activity of work. *Lancet*. 1953;265(6795): 1053–1057. contd.
 15. Li J, Siegrist J. Physical activity and risk of cardiovascular disease—a meta-analysis of prospective cohort studies. *Int J Environ Res Public Health*. 2012;9(2):391–407.
 16. Pescatello LS, MacDonald HV, Lamberti L, et al. Exercise for hypertension: a prescription update integrating existing recommendations with emerging research. *Curr Hypertens Rep*. 2015;17(11):87.
 17. Rubal BJ, Rosentswieg J, Hamerly B. Echocardiographic examination of women collegiate softball champions. *Med Sci Sports Exerc*. 1981;13(3):176–179.
 18. Wieling W, Borghols EA, Hollander AP, et al. Echocardiographic dimensions and maximal oxygen uptake in oarsmen during training. *Br Heart J*. 1981;46(2):190–195.
 19. Crouse SF, Rohack JJ, Jacobsen DJ. Cardiac structure and function in women basketball athletes: seasonal variation and comparisons with nonathletic controls. *Res Q Exerc Sport*. 1992;63(4):393–401.
 20. Pelliccia A, Maron BJ, Culasso F, et al. Athlete's heart in women. Echocardiographic characterization of highly trained elite female athletes. *JAMA*. 1996;276(3):211–215.
 21. Whyte G, Sharma S, George K, et al. Alterations in cardiac morphology and function in elite multi-disciplinary athletes. *Int J Sports Med*. 1999;20(4):222–226.
 22. Dzudie A, Menanga A, Hamadou B, et al. Ultrasonographic study of left ventricular function at rest in a group of highly trained black African handball players. *Eur J Echocardiogr*. 2007;8(2):122–127.
 23. Babaei Bigi MA, Aslani A. Aortic root size and prevalence of aortic regurgitation in elite strength trained athletes. *Am J Cardiol*. 2007;100(3):528–530.
 24. Carlsson L, Lind B, Laaksonen MS, et al. Enhanced systolic myocardial function in elite endurance athletes during combined arm-and-leg exercise. *Eur J Appl Physiol*. 2011;111(6):905–913.
 25. Caselli S, Di Pietro R, Di Paolo FM, et al. Left ventricular systolic performance is improved in elite athletes. *Eur J Echocardiogr*. 2011;12(7):514–519.
 26. Krol W, Braksator W, Kasprzak JD, et al. The influence of extreme mixed exertion load on the right ventricular dimensions and function in elite athletes: a tissue Doppler study. *Echocardiography*. 2011;28(7):753–760.
 - **This study reports no difference in aortic root diameter between athletes and controls after correction for Body Surface Area.**
 27. D'Andrea A, Cocchia R, Riegler L, et al. Aortic stiffness and distensibility in top-level athletes. *J Am Soc Echocardiogr*. 2012;25(5):561–567.
 28. Aparci M, Erdal M, Isilak Z, et al. Enlargement of the aorta: an occupational disease? *Exp Clin Cardiol*. 2013;18(2):93–97.
 29. Pelliccia A, Di Paolo FM, De Blasis E, et al. Prevalence and clinical significance of aortic root dilation in highly trained competitive athletes. *Circulation*. 2010;122(7):698–U621.
 30. Kinoshita N, Mimura J, Obayashi C, et al. Aortic root dilatation among young competitive athletes: echocardiographic screening of 1929 athletes between 15 and 34 years of age. *Am Heart J*. 2000;139(4):723–728.
 31. D'Andrea A, Cocchia R, Riegler L, et al. Aortic root dimensions in elite athletes. *Am J Cardiol*. 2010;105(11):1629–1634.
 32. Davis CK, Dyar DA, Vargas LA, et al. Cardiovascular and musculoskeletal assessment of elite US volleyball players. *Clin J Sport Med*. 2015;25(6):546–550.
 33. Bain MA, Zumwalt RE, Van der Bel-Kahn J. Marfan syndrome presenting as aortic rupture in a young athlete: sudden unexpected death? *Am J Forensic Med Pathol*. 1987;8(4):334–337.
 34. Schor JS, Horowitz MD, Livingstone AS. Recreational weight lifting and aortic dissection: case report review. *J Vasc Surg*. 1993;17(4):774–776.
 35. Baumgartner FJ, Omari BO, Robertson JM. Weight lifting, Marfan's syndrome, and acute aortic dissection. *Ann Thorac Surg*. 1997;64(6):1871–1872.
 36. Eleftheriades JA, Hatzaras I, Tranquilli MA, et al. Weight lifting and rupture of silent aortic aneurysms. *JAMA*. 2003;290(21):2803.
 37. Hatzaras I, Tranquilli M, Coady M, et al. Weight lifting and aortic dissection: more evidence for a connection. *Cardiology*. 2007;107(2):103–106.
 38. Uchida K, Imoto K, Yanagi H, et al. Acute aortic dissection occurring during the butterfly stroke in a 12-year-old boy. *Interact Cardiovasc Thorac Surg*. 2009;9(2):366–367.
 39. de Virgilio C, Nelson RJ, Milliken J, et al. Ascending aortic dissection in weight lifters with cystic medial degeneration. *Ann Thorac Surg*. 1990;49(4):638–642.
 40. Hogan CJ. An aortic dissection in a young weightlifter with non-Marfan fibrillinopathy. *Emerg Med J*. 2005;22(4):304–305.
 41. Ozyildirim S, Simsek EC, Gonencer M, et al. Acute aortic dissection and death owing to weight lifting in a patient presenting with chest pain. *Am J Cardiol*. 2015;115:5159–5160.
 42. Westaby S, Bradlow WM, Newton JD, et al. Rupture of an aneurysmal aortic diverticulum associated with coarctation and bicuspid aortic valve. *Circulation*. 2011;123(1):102–103.
 43. Chattranukulchai P, Luengtaviboon K, Tumkosit M. Tearing while bowling. *Case Reports*. 2013;2013:bcr2013010389.
 44. Cereda A, Sormani P, Russo CF, et al. Giant apparently not dissecting aneurysm and severe aortic valve insufficiency in an elite athlete. *Eur Heart J*. 2016;37(32):2527.
 45. Itagaki R, Kimura N, Itoh S, et al. Acute type a aortic dissection associated with a sporting activity. *Surg Today*. 2017;47(9):1163–1171.
 - **This study reports the amount of sports-related AD-A and focussed on the different types of sports practised during AD-A.**
 46. Gansera L, Deutsch O, Szameitat L, et al. Aortic dissections type a during sexual intercourse in male patients: accident or systematic coincidence? examination of 365 patients with acute aortic dissection within 20 years. *Thorac Cardiovasc Surg*. 2015;64(2):133–136.
 47. Schachner T, Fischler N, Dumfarth J, et al. Aortic dissection type A in alpine skiers. *Biomed Res Int*. 2013;192459:2013.
 48. Stefani L, Galanti G, Toncelli L, et al. Bicuspid aortic valve in competitive athletes. *Br J Sports Med*. 2008;42(1):31–35.
 49. Galanti G, Stefani L, Toncelli L, et al. Effects of sports activity in athletes with bicuspid aortic valve and mild aortic regurgitation. *Br J Sports Med*. 2010;44(4):275–279.
 - **This study reports aortic diameter growth rates in athletes with bicuspid and tricuspid aortic valve.**
 50. Stefani L, Galanti G, Innocenti G, et al. Exercise training in athletes with bicuspid aortic valve does not result in increased dimensions and impaired performance of the left ventricle. *Cardiol Res Pract*. 2014;2014.
 51. Spataro A, Pelliccia A, Rizzo M, et al. The natural course of bicuspid aortic valve in athletes. *Int J Sports Med*. 2008;29(1):81–85.
 52. Gibson C, Nielsen C, Alex R, et al. Mild aerobic exercise blocks elastin fiber fragmentation and aortic dilatation in a mouse model of Marfan syndrome associated aortic aneurysm. *J Appl Physiol*. 2017;123(1):147–160.
 53. Mas-Stachurska A, Siegert AM, Battle M, et al. Cardiovascular benefits of moderate exercise training in Marfan syndrome: insights from an animal model. *J Am Heart Assoc*. 2017;6:9.

- **This study reports a positive effect of mild-moderate dynamic exercise in mice with MFS.**
- 54. Benninghoven D, Hamann D, von Kodolitsch Y, et al. Inpatient rehabilitation for adult patients with Marfan syndrome: an observational pilot study. *Orphanet J Rare Dis.* 2017;12(1):127.
- 55. Iskandar A, Thompson PD. A meta-analysis of aortic root size in elite athletes. *Circulation.* 2013;127(7):791–798.
- 56. Wolak A, Gransar H, Thomson LE, et al. Aortic size assessment by noncontrast cardiac computed tomography: normal limits by age, gender, and body surface area. *JACC Cardiovasc Imaging.* 2008;1(2):200–209.
- 57. van der Werf C, van Langen IM, Wilde AA. Sudden death in the young: what do we know about it and how to prevent? *Circ Arrhythm Electrophysiol.* 2010;3(1):96–104.
- 58. Maron BJ. Sudden death in young athletes. *N Engl J Med.* 2003;349(11):1064–1075.
- 59. Maron BJ, Carney KP, Lever HM, et al. Relationship of race to sudden cardiac death in competitive athletes with hypertrophic cardiomyopathy. *J Am Coll Cardiol.* 2003;41(6):974–980.
- 60. Thompson PD, Franklin BA, Balady GJ, et al. Exercise and acute cardiovascular events placing the risks into perspective: a scientific statement from the American heart association council on nutrition, physical activity, and metabolism and the council on clinical cardiology. *Circulation.* 2007;115(17):2358–2368.
- 61. Mehta RH, Suzuki T, Hagan PG, et al. Predicting death in patients with acute type a aortic dissection. *Circulation.* 2002;105(2):200–206.
- 62. Ferencik M, Pape LA. Changes in size of ascending aorta and aortic valve function with time in patients with congenitally bicuspid aortic valves. *Am J Cardiol.* 2003;92(1):43–46.
- 63. La Canna G, Ficarra E, Tsagalau E, et al. Progression rate of ascending aortic dilation in patients with normally functioning bicuspid and tricuspid aortic valves. *Am J Cardiol.* 2006;98(2):249–253.
- 64. Detaint D, Michelena HI, Nkomo VT, et al. Aortic dilatation patterns and rates in adults with bicuspid aortic valves: a comparative study with Marfan syndrome and degenerative aortopathy. *Heart.* 2014;100(2):126–134.
- 65. Poulsen L. Animal models used to explore abdominal aortic aneurysms: a systematic review. *Eur J Vasc Endovasc Surg.* 2016;52(4):487–499.
- 66. Myers J, McElrath M, Jaffe A, et al. A randomized trial of exercise training in abdominal aortic aneurysm disease. *Med Sci Sports Exerc.* 2014;46(1):2–9.
- 67. Committee PAGA. 2018 Physical activity guidelines advisory committee scientific report. Washington, DC: U.S. Department of Health and Human Services; 2018.
- 68. Gezondheidsraad. Beweegrichtlijnen 2017. Den Haag: Gezondheidsraad; 2017. (publicatienr. /08).
- 69. Global Recommendations on Physical Activity for Health. Geneva: World Health Organization; 2010.
- 70. Olsson C, Franco-Cereceda A. Health-related quality of life in thoracic aortic disease: part I. Cases managed non-operatively. *Aorta.* 2013;1(3):153–161. *Stamford*